

Well-posedness of the Green-Naghdi and Boussinesq-Peregrine systems

VINCENT DUCHÊNE
SAMER ISRAWI

ABSTRACT. In this paper we address the Cauchy problem for two systems modeling the propagation of long gravity waves in a layer of homogeneous, incompressible and inviscid fluid delimited above by a free surface, and below by a non-necessarily flat rigid bottom. Concerning the Green-Naghdi system, we improve the result of Alvarez-Samaniego and Lannes [5] in the sense that much less regular data are allowed, and no loss of derivatives is involved. Concerning the Boussinesq-Peregrine system, we improve the lower bound on the time of existence provided by Mésognon-Gireau [42]. The main ingredient is a physically motivated change of unknowns revealing the quasilinear structure of the systems, from which energy methods are implemented.

1. Introduction

1.1. Motivation

The Green-Naghdi system¹ (sometimes called Serre or fully nonlinear Boussinesq system) is a model for the propagation of surface gravity waves in a layer of homogeneous incompressible inviscid fluid with rigid bottom and free surface. It has been formally derived several times in the literature, in particular in [48, 51, 25, 40, 47], using different techniques and various

VD is partially supported by the project Dyficolti ANR-13-BS01-0003-01 of the Agence Nationale de la Recherche.

SI is partially supported by the Lebanese University research program (MAA group project).

Keywords: Well-posedness theory, shallow water models, quasilinear dispersive systems.

Math. classification: 35L45, 35Q35, 76B15.

¹The Boussinesq-Peregrine system can be viewed as a simplification of the Green-Naghdi system for small-amplitude waves, which is particularly relevant for numerical purposes; see [45, 32, 42, 7]. This work is dedicated to the Green-Naghdi system and we only remark incidentally that our strategy may also be favorably applied to the Boussinesq-Peregrine system described thereafter.

hypotheses. For a clear and modern exposition, it is shown in [33] that the Green-Naghdi system can be derived as an asymptotic model from the water waves system (namely the “exact” equations for the propagation of surface gravity waves), by assuming that the typical horizontal length of the flow is much larger than the depth of the fluid layer—that is in the shallow-water regime—and that the flow is irrotational. Roughly speaking, a Taylor expansion with respect to the small “shallow-water parameter” yields at first order the Saint-Venant system, and at second order the Green-Naghdi system (see for instance [39] for higher order systems). As a relatively simple fully nonlinear model (that is without restriction on the amplitude of the waves) formally improving the precision of the Saint-Venant system, the Green-Naghdi system is widely used to model and numerically simulate the propagation of surface waves, in particular in coastal oceanography. It would be impossible to review the vast literature on the subject, and we only let the reader refer to [6, 10] for an introduction and relevant references.

In this work, we are interested in the structural properties and rigorous justification of the Green-Naghdi system. The derivation through formal Taylor expansions of Bonneton and Lannes [33] can be made rigorous [32, Prop. 5.8]: roughly speaking, any sufficiently smooth solution of the water waves system satisfies the Green-Naghdi system up to a quantifiable (small) remainder. This consistency result is only one step towards the full justification of the model in the following sense: the solution of the water waves system and the solution of the Green-Naghdi system with corresponding initial data remain close on a relevant time interval. In order for such result to hold, one needs of course to ensure the existence and uniqueness of a solution to the Green-Naghdi system in the aforementioned time interval for a large class of initial data; one also needs a stability property ensuring that the two solutions are close. These two results typically call for robust energy estimates on exact and approximate solutions.

Somewhat surprisingly, the well-posedness theory concerning the Cauchy problem for the Green-Naghdi system is in some sense less satisfactory than the corresponding one for the water waves system. Again, the literature on the latter problem is too vast to summarize, and we only mention the result of [4, 27] and [32, Theorem 4.16]. Indeed, the latter pay attention to the various dimensionless parameters of the system, and in particular obtain results which hold uniformly with respect to the shallow-water parameter. The outcome of these results is that provided that the

initial data is sufficiently regular (measured through Sobolev spaces) and satisfy physical assumptions —the so-called non-cavitation and Rayleigh-Taylor criteria— then there exists a unique solution of the water waves system preserving the regularity of the initial data. Moreover, the maximal time of existence may be bounded from below uniformly with respect to the shallow-water parameter; see details therein. Such result is very much nontrivial as the limit of small shallow-water parameter is singular in some sense. Similar results have been proved for the Green-Naghdi system in horizontal dimension $d = 1$ in [38] (for flat bottom) and [29] (for general bathymetries). An apparent obstruction prevents to implement the strategy in dimension $d = 2$; see the discussion therein. Alvarez-Samaniego and Lannes [5] proved an existence and uniqueness result in the general setting and on the correct time-scale but their proof relies on a Nash-Moser scheme, and as such involves a loss of derivatives between the regularity of the initial data and the control of the solution at positive times. *The main result of this paper is to show that this loss of derivatives is in fact not necessary, and that the Cauchy problem for the Green-Naghdi system is well-posed in the sense of Hadamard in Sobolev-type spaces.*

1.2. Strategy

Let us now introduce the system of equations at stake. In order to ease the discussion and notations, we restrict the study to the horizontal space $X \in \mathbb{R}^d$ with $d = 2$, although the results are easily adapted to the setting $d = 1$, thus yielding another proof of the result in [29]. The non-dimensionalized Green-Naghdi system may be written (see [33, 32])

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ (\text{Id} + \mu\mathcal{T}[h, \beta b])\partial_t \mathbf{u} + \nabla \zeta + \epsilon(\mathbf{u} \cdot \nabla)\mathbf{u} + \mu\epsilon(\mathcal{Q}[h, \mathbf{u}] + \mathcal{Q}_b[h, \beta b, \mathbf{u}]) = 0, \end{cases} \quad (1.1)$$

with $h = 1 + \epsilon\zeta - \beta b$ and

$$\begin{aligned} \mathcal{T}[h, \beta b]\mathbf{u} \stackrel{\text{def}}{=} & \frac{-1}{3h}\nabla(h^3\nabla \cdot \mathbf{u}) + \frac{1}{2h}\left(\nabla(h^2(\beta\nabla b) \cdot \mathbf{u}) - h^2(\beta\nabla b)\nabla \cdot \mathbf{u}\right) \\ & + \beta^2(\nabla b \cdot \mathbf{u})\nabla b, \quad (1.2) \end{aligned}$$

and

$$\begin{aligned} \mathcal{Q}[h, \mathbf{u}] &\stackrel{\text{def}}{=} \frac{-1}{3h} \nabla \left(h^3 ((\mathbf{u} \cdot \nabla)(\nabla \cdot \mathbf{u}) - (\nabla \cdot \mathbf{u})^2) \right), \\ \mathcal{Q}_b[h, \beta b, \mathbf{u}] &\stackrel{\text{def}}{=} \frac{\beta}{2h} \left(\nabla (h^2 (\mathbf{u} \cdot \nabla)^2 b) - h^2 ((\mathbf{u} \cdot \nabla)(\nabla \cdot \mathbf{u}) - (\nabla \cdot \mathbf{u})^2) \nabla b \right) \\ &\quad + \beta^2 ((\mathbf{u} \cdot \nabla)^2 b) \nabla b. \end{aligned}$$

Here, the unknowns are $\zeta(t, X) \in \mathbb{R}$ and $\mathbf{u}(t, X) \in \mathbb{R}^d$ (representing respectively the dimensionless surface deformation and layer-averaged horizontal velocity), $b(X) \in \mathbb{R}$ is the fixed bottom topography (so that $h(t, X)$ represents the depth of the fluid layer) and ϵ, β, μ are dimensionless parameters.

As aforementioned, by setting $\mu = 0$ in (1.1), one recovers the Saint-Venant system, which is an archetype of first-order quasilinear systems of conservation laws. Our strategy in the following is to adapt to the Green-Naghdi equations the well-known techniques — and in particular *a priori* energy estimates — developed for such systems. Such energy estimates are obviously not guaranteed, due to the presence of the additional third-order nonlinear operators. The key ingredient of this work is the extraction of a quasilinear structure of (1.1), from which energy estimates can be deduced, and eventually a standard Picard iteration scheme can be set up.

Such a “quasilinearization” is also the key ingredient in the proof of the local existence of solutions to the water waves system [32, Theorem 4.16]. However, the structure of the water waves system and the one of the Green-Naghdi system look different, due to the fact that they use different unknowns. Indeed, the second equation in (1.1) describes the time-evolution of the layer-averaged horizontal velocity, while the Zakharov/Craig-Sulem formulation of the water waves system involves the trace of the velocity potential at the surface. To our opinion, the main contribution of this work is the demonstration that when expressed in a different set of variables, the Green-Naghdi system possesses a structure which is very similar to the water waves one; and that one can take advantage of this fact to adapt the proof of the local well-posedness of the latter to the one of the former.

To be more precise, we work on another formulation for system (1.1):

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h \mathbf{u}) = 0, \\ (\partial_t + \epsilon \mathbf{u}^\perp \text{curl}) \mathbf{v} + \nabla \zeta + \frac{\epsilon}{2} \nabla (|\mathbf{u}|^2) = \mu \epsilon \nabla (\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]), \end{cases} \quad (1.3)$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

where we denote $\text{curl}(v_1, v_2) \stackrel{\text{def}}{=} \partial_1 v_2 - \partial_2 v_1$ and $(u_1, u_2)^\perp \stackrel{\text{def}}{=} (-u_2, u_1)$,

$$\mathcal{R}[h, \mathbf{u}] \stackrel{\text{def}}{=} \frac{\mathbf{u}}{3h} \cdot \nabla(h^3 \nabla \cdot \mathbf{u}) + \frac{1}{2} h^2 (\nabla \cdot \mathbf{u})^2, \quad (1.4)$$

$$\mathcal{R}_b[h, \beta b, \mathbf{u}] \stackrel{\text{def}}{=} -\frac{1}{2} \left(\frac{\mathbf{u}}{h} \cdot \nabla(h^2(\beta \nabla b \cdot \mathbf{u})) + h(\beta \nabla b \cdot \mathbf{u}) \nabla \cdot \mathbf{u} + (\beta \nabla b \cdot \mathbf{u})^2 \right) \quad (1.5)$$

and \mathbf{v} is defined (recalling the definition of the operator \mathcal{T} in (1.2)) by

$$h\mathbf{v} = h\mathbf{u} + \mu h \mathcal{T}[h, \beta b] \mathbf{u} \stackrel{\text{def}}{=} \mathfrak{I}[h, \beta b] \mathbf{u}. \quad (1.6)$$

In the following, we study system (1.3) as evolution equations for the variables ζ and \mathbf{v} , with $\mathbf{u} = \mathbf{u}[h, \beta b, \mathbf{v}]$ being uniquely defined (see Lemma 2.3 thereafter) by (1.6), and deduce the well-posedness of (1.1) from the one of (1.3).

That system (1.3) is equivalent to (1.1) is certainly not straightforward, and we detail the calculations in Section 6. Physically speaking, the variable \mathbf{v} approximates (in the shallow-water regime, $\mu \ll 1$) $\mathbf{v}_{\text{ww}} = \nabla \psi_{\text{ww}}$ where ψ_{ww} is the trace of the velocity potential at the surface, and thus system (1.3) is more directly comparable to the water waves system. In particular, the Hamiltonian structure of system (1.3), as brought to light in [26, 37], is a direct counterpart of the celebrated one of the water waves system, and we show in Appendix B how system (1.3) can be quickly derived thanks to the Hamiltonian formalism. This allows to obtain preserved quantities of the system in a straightforward way (see [50] and references therein). Most importantly for our purposes, this allows us to follow the strategy of the proof for the local existence of a solution to the water waves system in [27, 32], and to obtain the corresponding local existence result for system (1.3). More precisely, the variables ζ, \mathbf{v} allow to define the analogue of Alinhac’s “good unknowns” [3] for the water waves system (see *e.g.* [2]) on which energy estimates can be established. We then extend the analysis so as to prove the well-posedness of the Cauchy problem, in the sense of Hadamard.

After the completion of this work, it was pointed to us that the use of system (1.3) was not necessary (and may be an over-complicated strategy) to derive *a priori* energy estimates. For the sake of completeness, we sketch in Appendix A the computations which provide such *a priori* estimates directly on system (1.1) and would yield an alternative proof of our main results. As a matter of fact, a sketch of such proof was given in [21].

However, we believe that the similarity of structure between the Green-Naghdi system and the one of the water waves system that we exhibit and exploit in this work is an interesting feature. It may serve as a pedagogical tool to get a grasp at some properties of the latter without technical difficulties related to the Dirichlet-to-Neumann operator. Incidentally, we do not claim the discovery of formulation (1.3); see [35] and references given therein, as well as in Appendix B.

1.3. Main results

Let us now present the main results of this work. Here and thereafter, we fix the parameters $\epsilon, \beta \geq 0$ and $\mu \in (0, \mu^*)$ with $\mu^* > 0$. The validity of the Green-Naghdi system as an asymptotic model for the water waves system stems from assuming $\mu \ll 1$ while $\epsilon, \beta = \mathcal{O}(1)$ —see [32] for details— but we do not make use of such restriction in this work. However, we shall always assume that

$$0 < h_* < h(\epsilon\zeta(x), \beta b(x)) < h^* < \infty, \quad h(\epsilon\zeta, \beta b) \stackrel{\text{def}}{=} 1 + \epsilon\zeta - \beta b. \quad (1.7)$$

We work with the following functional spaces, defined for $n \in \mathbb{N}$ by

$$\begin{aligned} H^n &\stackrel{\text{def}}{=} \{\zeta \in L^2(\mathbb{R}^d), |\zeta|_{H^n}^2 \stackrel{\text{def}}{=} \sum_{|\alpha|=0}^n |\partial^\alpha \mathbf{u}|_{L^2}^2 < \infty\}, \\ \dot{H}^n &\stackrel{\text{def}}{=} \{b \in L^2_{\text{loc}}(\mathbb{R}^d), \nabla b \in (H^{n-1})^d\}, \\ X^n &\stackrel{\text{def}}{=} \{\mathbf{u} \in L^2(\mathbb{R}^d)^d, |\mathbf{u}|_{X^n}^2 \stackrel{\text{def}}{=} \sum_{|\alpha|=0}^n |\partial^\alpha \mathbf{u}|_{L^2}^2 + \mu |\partial^\alpha \nabla \cdot \mathbf{u}|_{L^2}^2 < \infty\}, \\ Y^n &\stackrel{\text{def}}{=} \{\mathbf{v} \in (X^0)', |\mathbf{v}|_{Y^n}^2 \stackrel{\text{def}}{=} \sum_{|\alpha|=0}^n |\partial^\alpha \mathbf{v}|_{(X^0)'}^2 < \infty\}. \end{aligned}$$

Here, $\alpha \in \mathbb{N}^d$ is a multi-index, $(X^0)'$ is the topological dual space of X^0 , endowed with the norm of the strong topology; and we denote by $\langle \mathbf{v}, \mathbf{u} \rangle_{(X^0)'}$ the $(X^0)' - X^0$ duality bracket.

Given variables $\lambda_i \in \mathbb{R}^+$, we denote $C(\lambda_1, \lambda_2, \dots)$ a multivariate polynomial with non-negative coefficients, and $F(\lambda_1, \lambda_2, \dots)$ a multivariate polynomial with non-negative coefficients and zero constant term. Since such notations are used for upper bounds and will take variables restricted

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

to line segments, C should be regarded as a constant and F as a linear functional.

Theorem 1.1 (Well-posedness of the Green-Naghdi system). *Let $N \geq 4$, $b \in \dot{H}^{N+2}$ and $(\zeta_0, \mathbf{u}_0) \in H^N \times X^N$ satisfying (1.7) with $h_\star, h^\star > 0$. Then there exists $T > 0$ and a unique $(\zeta, \mathbf{u}) \in C([0, T]; H^N \times X^N)$ satisfying (1.1) with initial data $(\zeta, \mathbf{u})|_{t=0} = (\zeta_0, \mathbf{u}_0)$. Moreover, one can restrict*

$$T^{-1} = C(\mu^\star, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^N}, \epsilon|\mathbf{u}_0|_{X^N}) > 0$$

such that, for any $t \in [0, T]$, (1.7) holds with $\tilde{h}_\star = h_\star/2$, $\tilde{h}^\star = 2h^\star$, and

$$\sup_{t \in [0, T]} (|\zeta|_{H^N}^2 + |\mathbf{u}|_{X^N}^2) \leq \mathbf{C} \times (|\zeta_0|_{H^N}^2 + |\mathbf{u}_0|_{X^N}^2),$$

where $\mathbf{C} = C(\mu^\star, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^N}, \epsilon|\mathbf{u}_0|_{X^N})$, and such that the map $(\zeta_0, \mathbf{u}_0) \in H^N \times X^N \mapsto (\zeta, \mathbf{u}) \in C([0, T]; H^N \times X^N)$ is continuous.

The full justification of the Green-Naghdi system is a consequence of the local existence and uniqueness result for the water waves system ([32, Th. 4.16]), its consistency with the Green-Naghdi system ([32, Prop. 5.8]), the well-posedness of the Cauchy problem for the Green-Naghdi system (Theorem 1.1) as well as a stability result ensuring the Lipschitz dependence of the error with respect to perturbations of the system (Proposition 5.2, thereafter). The following result improves [32, Th. 6.15] by the level of regularity required on the initial data.

Theorem 1.2 (Full justification of the Green-Naghdi system). *Let $N \geq 7$ and assume that $(\zeta_{\text{ww}}, \psi_{\text{ww}}) \in C([0, T_{\text{ww}}]; H^N \times \dot{H}^{N+1})$ is a solution to the water waves system (B.3) (recall that such solutions exist for a large class of initial data by [32, Th. 4.16]) such that (1.7) holds with $b \in H^{N+2}(\mathbb{R}^d)$.*

Denote $\zeta_0 = (\zeta_{\text{ww}})|_{t=0}$ and $h_0 = 1 + \epsilon\zeta_0 - \beta b$, $\psi_0 = (\psi_{\text{ww}})|_{t=0}$ and $\mathbf{u}_0 = \mathfrak{T}[h_0, \beta b]^{-1}(h_0 \nabla \psi_0)$; and $M = \sup_{t \in [0, T_{\text{ww}}]} (|\zeta_{\text{ww}}|_{H^N} + |\nabla \psi_{\text{ww}}|_{H^N})$.

Then there exists $T > 0$ and $(\zeta_{\text{GN}}, \mathbf{u}_{\text{GN}}) \in C([0, T]; H^N \times X^N)$ unique strong solution to the Green-Naghdi system (1.1) with initial data (ζ_0, \mathbf{u}_0) , by Theorem 1.1; and one can restrict

$$T^{-1} = C(\mu^\star, h_\star^{-1}, h^\star)F(\beta|b|_{H^{N+2}}, \epsilon M) > 0$$

such that for any $t \in [0, \min(T, T_{\text{ww}})]$,

$$|\zeta_{\text{ww}} - \zeta_{\text{GN}}|_{H^{N-6}} + |\nabla \psi_{\text{ww}} - \mathbf{v}_{\text{GN}}|_{Y^{N-6}} \leq \mathbf{C} \mu^2 t,$$

with $\mathbf{C} = C(\mu^\star, h_\star^{-1}, h^\star, \beta|b|_{H^{N+2}}, \epsilon M)$, and \mathbf{v}_{GN} is defined by (1.6).

Remark 1.3. The functional spaces arise as natural energy spaces for the quasilinear structure of system (1.3) that we exhibit in this work. The regularity assumption $N \geq 4$ is most certainly not optimal. By the method of our proof, we are restricted to $N \in \mathbb{N}$, and we cannot hope to obtain a lower threshold than in the pure quasilinear setting, namely $N > d/2 + 1$. That $N = 3$ is not allowed when $d = 2$ comes from technical limitations in various places.

Since our proof is based solely on the structural properties of the system and on energy estimates (in particular, no dispersive estimates are used), it may be adapted almost verbatim to the periodic situation. Although we have not checked all the technical details, we also expect that the strategy may be extended to the more general situation of Kato’s uniformly local Sobolev spaces [30].

Remark 1.4. Applying the operator curl to (1.3)₂, one observes the identity

$$\partial_t \operatorname{curl} \mathbf{v} + \epsilon \nabla \cdot (\mathbf{u} \operatorname{curl} \mathbf{v}) = 0.$$

Thus standard energy estimates on conservation laws [8] yield, for any $t \in [0, T]$,

$$|\operatorname{curl} \mathbf{v}|_{H^{N-1}}(t) \leq |\operatorname{curl} \mathbf{v}_0|_{H^{N-1}} \exp(\epsilon \lambda t)$$

with $\lambda \lesssim |\mathbf{u}|_{H^N} \leq |\mathbf{u}|_{X^N}$. Thus a byproduct of Theorem 1.1 is that initial smallness of the “generalized vorticity” [40, 24], $\operatorname{curl} \mathbf{v}$, propagates for large positive times —and remains trivial if it vanishes initially, as in Theorem 1.2. As already mentioned, the variable \mathbf{v} physically approximates $\nabla \psi_{\text{ww}}$ where ψ_{ww} is the trace of the velocity potential at the surface. It is therefore physically relevant to assume

$$\operatorname{curl} \mathbf{v} = \operatorname{curl} (\mathbf{u} + \mu \mathcal{T}[h, \beta b] \mathbf{u}) \equiv 0.$$

This two-dimensional irrotationality condition is a direct consequence of the three-dimensional irrotationality assumption on the velocity flow inside the fluid layer. Outside of this irrotational framework, we believe that the Green-Naghdi system is not a valid model, and refer to [14] for a thorough discussion in this situation.

Outline. The body of the paper is dedicated to the proof of Theorem 1.1. In Section 2, we provide some technical results concerning our functional spaces and the operator \mathfrak{T} . We then exhibit the quasilinear system satisfied by the derivatives of any regular solution to (1.3) in Section 3.

This quasilinear formulation allows to obtain *a priori* energy estimates in Section 4. Finally, we make use of these energy estimates for proving the well-posedness of the Cauchy problem for system (1.3) in Section 5. The equivalence between formulation (1.3) and (1.1) is stated in Proposition 6.1. Theorems 1.1 and 1.2 follow as direct consequence of the results in Section 5 and Proposition 6.1, and we complete the proofs in Section 6. In Appendix A, we roughly sketch of how energy estimates could be obtained directly on formulation (1.1). We also provide a quick discussion on the derivation and Hamiltonian formulation of system (1.3) in Appendix B. We conclude this section with some independent remarks.

Large time well-posedness and asymptotics. A very natural question in the oceanographic context concerns the large time asymptotic behavior of solutions to the Green-Naghdi system for small data. After a straightforward rescaling of (1.1), the problem is naturally formulated in terms of solutions to the system

$$\begin{cases} \partial_t \zeta + \frac{1}{\epsilon} \nabla \cdot (h \mathbf{u}) = 0, & h \stackrel{\text{def}}{=} 1 + \epsilon \zeta - \beta b, \\ (\text{Id} + \mu \mathcal{T}[h, \beta b]) \partial_t \mathbf{u} + \frac{1}{\epsilon} \nabla \zeta + (\mathbf{u} \cdot \nabla) \mathbf{u} + \mu (\mathcal{Q}[h, \mathbf{u}] + \mathcal{Q}_b[h, \beta b, \mathbf{u}]) = 0. \end{cases} \quad (1.8)$$

Is the Cauchy problem for (1.8) locally well-posed, uniformly with respect to the small parameter ϵ ? Is it globally well-posed for ϵ small enough? Can we exhibit “averaged” equations asymptotically describing a slow coherent evolution of the solution?

This type of singular limit has been widely studied in particular in the context of the low Mach number limit; see *e.g.* [22, 1] and references therein. As a matter of fact, when $\beta = \mu = 0$, one recognizes the incompressible limit for the isentropic two-dimensional Euler equations, and it is tempting to elaborate on the analogy. One would then expect the solutions to (1.8) to be asymptotically described (as $\epsilon \rightarrow 0$) as the superposition of two components, described thereafter.

- (1) The “incompressible” component, being defined as the solution to

$$\begin{cases} \nabla \cdot ((1 - \beta b) \mathbf{u}) = 0, \\ (\text{Id} + \mu \mathcal{T}[1 - \beta b, \beta b]) \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} \\ \quad + \mu (\mathcal{Q}[1 - \beta b, \mathbf{u}] + \mathcal{Q}_b[1 - \beta b, \beta b, \mathbf{u}]) = -\nabla p, \end{cases} \quad (1.9)$$

where the “pressure” p in $(1.9)_2$ is the Lagrange multiplier associated to the “incompressibility” constraint $(1.9)_1$.

(2) The “acoustic” component, being defined as the solution to

$$\begin{cases} \partial_t \zeta + \frac{1}{\epsilon} \nabla \cdot ((1 - \beta b) \mathbf{u}) = 0, \\ (\text{Id} + \mu \mathcal{T}[1 - \beta b, \beta b]) \partial_t \mathbf{u} + \frac{1}{\epsilon} \nabla \zeta = 0, \end{cases} \quad (1.10)$$

with initial data satisfying $\text{curl}(\mathbf{u} + \mu \mathcal{T}[1 - \beta b, \beta b] \mathbf{u})|_{t=0} = 0$.

System (1.9) was derived in [12, 13], and is usually referred to as the *great-lake equations*. Its well-posedness, extending the theory concerning the two-dimensional incompressible Euler equations, was subsequently provided in [36, 44]. However, these authors apparently overlooked the role of the irrotationality assumption discussed in Remark 1.4, as the only functions satisfying the constraint $(1.9)_1$ as well as the irrotationality condition $\text{curl}(\mathbf{u} + \mu \mathcal{T}[1 - \beta b, \beta b] \mathbf{u}) = 0$ are in fact trivial. In other words, in the irrotational framework that is the only one so far for which the Green-Naghdi system is rigorously justified, the incompressible (or rigid-lid) component vanishes; see also [43] for a similar discussion on the water waves system. One thus expects that the flow is asymptotically described by (1.10) only, in the limit $\epsilon \rightarrow 0$ and in the irrotational setting.

However, when trying to adapt the usual strategy for rigorously proving such behavior, one immediately encounters a serious difficulty in the physically relevant situation of non-trivial topography, which transpires in the the fact that our lower bound for the existence time in Theorem 1.1 depends on the size of the bottom variations in addition to the size of the initial data. When transcribed to system (1.8), this means that we are not able to obtain a lower bound on the existence time of its solutions which is uniform with respect to ϵ , unless $\beta = \mathcal{O}(\epsilon)$.

For the Saint-Venant system, that is setting $\mu = 0$, Bresch and Métivier [11] have obtained such a uniform lower bound without any restriction on the amplitude bathymetry. The strategy consists in estimating first the time derivatives of the solution, and then using the system to deduce estimates on space derivatives. A related strategy (in the sense that we look for operators commuting with the singular component of the system) amounts to remark that for any $n \in \mathbb{N}$, one can control the L^2 -norm of

$$\zeta_n \stackrel{\text{def}}{=} (\nabla \cdot (1 - b) \nabla)^n \zeta, \quad \mathbf{u}_n \stackrel{\text{def}}{=} (\nabla(1 - b) \nabla \cdot)^n \mathbf{u}$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

by exhibiting the quasilinear system satisfied by (ζ_n, \mathbf{u}_n) and applying simple energy estimates. This allows to control the H^{2n} -norm of ζ, \mathbf{u} , provided that the initial data and bottom topography are sufficiently regular. One expects a similar strategy to work for the water waves system (partial results have been obtained by the method of time derivatives in [41]), that is to control

$$\zeta_n \stackrel{\text{def}}{=} \left(\frac{1}{\mu} G^\mu[0, \beta b]\right)^n \zeta, \quad \psi_n \stackrel{\text{def}}{=} \left(\frac{1}{\mu} G^\mu[0, \beta b]\right)^n \psi$$

where G^μ is the Dirichlet-to-Neumann operator, recalled in Appendix B. Since G^μ is an operator of order 1, controlling ζ_n, ψ_n indeed allows to control higher regularities on ζ, ψ . The strategy however fails for the Green-Naghdi system, as the corresponding operator, namely (see Appendix B)

$$\frac{1}{\mu} G^\mu[0, \beta b] \bullet \approx -\nabla \cdot \left((1 - \beta b) \mathfrak{T}[1 - \beta b, \beta b]^{-1} \{ (1 - \beta b) \nabla \bullet \} \right)$$

is of order 0. One could easily propose different systems that do not suffer from such a shortcoming, by adding the effect of surface tension as in [41], or modifying the system without hurting its consistency as in [42]. This is however out of the scope of the present work, and we leave the question of uniform lower bounds for the existence time of solutions to (1.8) as an open problem.

Other models. We expect that our strategy may be of interest to other models for surface gravity waves. Since our result holds uniformly with respect to $\mu \in (0, 1)$, we may deduce the well-posedness of the Saint-Venant system. This result is of course a direct application of the standard theory on quasilinear hyperbolic systems [8]. In the other direction, it would be interesting to apply our strategy to the higher-order models derived by Matsuno [39], which enrich the Saint-Venant and Green-Naghdi systems with models of arbitrary high order, while preserving the structure of which we take advantage in this work. Similarly, one can derive models with improved frequency dispersion while preserving the structure of the Green-Naghdi model, by modifying the approximate Hamiltonian in Appendix B. Such a strategy was applied by the authors in the one-dimensional and bilayer situation in [20]. Interestingly, since such models can be tuned to fit the dispersion relation of the water waves system, they do not suffer from the shortcoming described above, so that large-time well-posedness is expected to hold even in the presence of a non-trivial bathymetry.

Let us clarify however that our strategy does not require the Hamiltonian structure, but only exploits the ability to construct “good unknowns” through ζ, \mathbf{v} and their derivatives. Thus even models which are derived through careless approximations may preserve the quasilinear structure which is necessary for our energy estimates. Such is the case for the Boussinesq-Peregrine system (see [45, 32, 42]), which consists in a simplification of the Green-Naghdi system obtained by withdrawing contributions of size $\mathcal{O}(\mu\epsilon)$ while keeping $\mathcal{O}(\mu\beta)$ ones:

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ \partial_t (\mathbf{u} + \mu\mathcal{T}[1 - \beta b, \beta b]\mathbf{u}) + \nabla \zeta + \epsilon(\mathbf{u} \cdot \nabla)\mathbf{u} = 0. \end{cases} \quad (1.11)$$

Because the elliptic operator $\text{Id} + \mu\mathcal{T}[1 - \beta b, \beta b]$ does not depend on time, solving numerically the Boussinesq-Peregrine system is much less costly than solving the Green-Naghdi system; see [35, 34]. It turns out that the proof of Theorem 1.1 does extend to the Boussinesq-Peregrine system with straightforward modifications (all on the side of simplification). More precisely, we obtain the well-posedness of system (1.11) after applying the change of variable

$$\mathbf{v} \stackrel{\text{def}}{=} \mathbf{u} + \mu\mathcal{T}[1 - \beta b, \beta b]\mathbf{u} \quad (1.12)$$

and extracting the quasilinear system satisfied by $\partial^\alpha \zeta$ and $\partial^\alpha \mathbf{v}$.

Theorem 1.5 (Well-posedness and full justification of the Boussinesq-Peregrine system). *Theorem 1.1 holds replacing system (1.1) with system (1.11). The full justification in the sense of Theorem 1.2 holds as well, controlling the error as*

$$|\zeta_{\text{ww}} - \zeta_{\text{BP}}|_{H^{N-6}} + |\nabla \psi_{\text{ww}} - \mathbf{v}_{\text{BP}}|_{Y^{N-6}} \leq \mathbf{C}(\mu^2 + \mu\epsilon)t,$$

with $\zeta_{\text{BP}}, \mathbf{u}_{\text{BP}}$ the unique strong solution to (1.11) and \mathbf{v}_{BP} defined by (1.12), and the other notations are as in Theorem 1.2.

Compared with [42, Th. 2.1], our result provides a larger lower bound on the time of existence, and does not rely on the assumption $\epsilon = \mathcal{O}(\mu)$.

2. Preliminary results

In this section, we fix parameters $n \in \mathbb{N}$, $\alpha \in \mathbb{N}^d$, and dimension $d \in \{1, 2\}$. We denote $a \lesssim b$ for $a \leq Cb$ where C is a constant depending (non-decreasingly) only on n, d , and possibly $|\alpha|$ and μ . We denote $\langle A \rangle_{n>r} = A$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

if $n > r$ and $\langle A \rangle_{n>r} = 0$ otherwise, and $a \vee b = \max(a, b)$. The results are tailored for the dimension $d = 2$ (through the repeated use of the continuous Sobolev embedding $H^2 \subset L^\infty$ for instance) but hold as well when $d = 1$. They are not meant to be sharp, but only sufficient for our needs. The call for “tame” estimates stems from the Bona-Smith technique enforcing continuity properties stated in Proposition 5.3; rougher estimates would be sufficient for the existence and uniqueness of solutions, as in Proposition 5.1.

Lemma 2.1. *The continuous embeddings $H^{n+1}(\mathbb{R}^d)^d \subset X^n \subset H^n(\mathbb{R}^d)^d$ and $H^n(\mathbb{R}^d)^d \subset Y^n \subset H^{n-1}(\mathbb{R}^d)^d$ hold. The following inequalities hold as soon as the right-hand side is finite:*

$$|\mathbf{u}|_{H^n} \leq |\mathbf{u}|_{X^n}, \quad |\mathbf{u}|_{X^n} \lesssim |\mathbf{u}|_{H^{n+1}}, \quad (2.1)$$

$$|\mathbf{v}|_{H^{n-1}} \lesssim |\mathbf{v}|_{Y^n}, \quad |\mathbf{v}|_{Y^n} \leq |\mathbf{v}|_{H^n}. \quad (2.2)$$

We also have the non-uniform continuous embedding

$$|\nabla f|_{Y^n} \lesssim \frac{1}{\sqrt{\mu}} |f|_{H^n}, \quad |\nabla \cdot \mathbf{u}|_{H^n} \lesssim \frac{1}{\sqrt{\mu}} |\mathbf{u}|_{X^n}. \quad (2.3)$$

Proof. The continuous embeddings $H^1(\mathbb{R}^d)^d \subset X^0 \subset L^2(\mathbb{R}^d)^d$ are straightforward, and the corresponding $L^2(\mathbb{R}^d)^d \subset Y^0 \subset H^{-1}(\mathbb{R}^d)^d$ follow by duality. The estimate (2.3) with $n = 0$ is easily checked, as for any $\mathbf{u} \in X^0$,

$$|\langle \nabla f, \mathbf{u} \rangle_{(X^0)'}| = |(f, \nabla \cdot \mathbf{u})_{L^2}| \leq \frac{1}{\sqrt{\mu}} |f|_{L^2} |\mathbf{u}|_{X^0}.$$

The case $n \in \mathbb{N}^*$ is reduced to the case $n = 0$ by considering $\partial^\alpha \mathbf{u}$, $\partial^\alpha \mathbf{v}$, $\partial^\alpha f$ with $0 \leq |\alpha| \leq n$. \square

Lemma 2.2. *Let $f \in H^{2\vee n}(\mathbb{R}^d)$ and $g \in H^n(\mathbb{R}^d)$. Then $fg \in H^n(\mathbb{R}^d)$*

$$|fg|_{H^n} \lesssim |f|_{H^2} |g|_{H^n} + \langle |f|_{H^n} |g|_{H^2} \rangle_{n>2}. \quad (2.4)$$

The above holds as well allowing exceptionally the value $n = -1$.

Let $f \in L^\infty(\mathbb{R}^d) \cap \dot{H}^{3\vee n}(\mathbb{R}^d)$ and $g \in H^n(\mathbb{R}^d)$. Then $fg \in H^n(\mathbb{R}^d)$ and

$$|fg|_{H^n} \lesssim (|f|_{L^\infty} + |\nabla f|_{H^2}) |g|_{H^n} + \langle |\nabla f|_{H^{n-1}} |g|_{H^2} \rangle_{n>2}. \quad (2.5)$$

If, moreover, $f \geq f_0 > 0$, then $f^{-1}g \in H^n(\mathbb{R}^d)$ and

$$|f^{-1}g|_{H^n} \leq C(f_0^{-1}, |f|_{L^\infty}, |\nabla f|_{H^2}) (|g|_{H^n} + \langle |\nabla f|_{H^{n-1}} |g|_{H^2} \rangle_{n>2}). \quad (2.6)$$

Let $f \in L^\infty(\mathbb{R}^d) \cap \dot{H}^{3\nu n+1}(\mathbb{R}^d)$ and $\mathbf{u} \in X^n$. Then $f\mathbf{u} \in X^n$ and

$$|f\mathbf{u}|_{X^n} \lesssim (|f|_{L^\infty} + |\nabla f|_{H^2})|\mathbf{u}|_{X^n} + \langle |\nabla f|_{H^n}|\mathbf{u}|_{X^2} \rangle_{n>2}. \quad (2.7)$$

Let $f \in L^\infty(\mathbb{R}^d) \cap \dot{H}^{3\nu n}(\mathbb{R}^d)$, and $\mathbf{v} \in Y^n$. One has $f\mathbf{v} \in Y^n$ and

$$|f\mathbf{v}|_{Y^n} \lesssim (|f|_{L^\infty} + |\nabla f|_{H^2})|\mathbf{v}|_{Y^n} + \langle |\nabla f|_{H^{n-1}}|\mathbf{v}|_{H^2} \rangle_{n>2}. \quad (2.8)$$

Proof. Estimate (2.4) is well-known; see [32, Prop. B.2] for instance. As for (2.5), the cases $n \in \{0, 1, 2\}$ are straightforward, using Leibniz rule and the continuous Sobolev embedding $H^2 \subset L^\infty$. When $n > 2$, we decompose Leibniz rule as follows: for any $1 \leq |\alpha| \leq n$,

$$\partial^\alpha(fg) = f\partial^\alpha g + \sum_{\substack{\beta+\gamma=\alpha \\ |\beta|\geq 1, |\gamma|\geq 0}} \binom{\alpha}{\beta} (\partial^\beta f)(\partial^\gamma g).$$

Since $|\beta| \geq 1$ and using the standard bilinear estimate for (see *e.g.* [52, Prop. 3.6]), one has

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \lesssim |\nabla f|_{L^\infty}|g|_{H^{|\beta|+|\gamma|-1}} + |\nabla f|_{H^{|\beta|+|\gamma|-1}}|g|_{L^\infty},$$

and the result follows. Estimate (2.6) is obtained in the same way, and using the induction hypothesis to control the contribution of $|\nabla(f^{-1})|_{H^{|\beta|+|\gamma|-1}}$. As for (2.7), we use

$$|f\mathbf{u}|_{X^n} \lesssim |f\mathbf{u}|_{H^n} + \sqrt{\mu}|f\nabla \cdot \mathbf{u}|_{H^n} + \sqrt{\mu}|\nabla f \cdot \mathbf{u}|_{H^n},$$

with product estimates (2.4) and (2.5), as well as the continuous embedding (2.1) in Lemma 2.1. Finally, we notice that $f\mathbf{v} \in Y^0 = (X^0)'$ and $|f\mathbf{v}|_{Y^0} \lesssim (|f|_{L^\infty} + |\nabla f|_{H^2})|\mathbf{v}|_{(X^0)'}$, since for any $\mathbf{u} \in X^0$,

$$|\langle f\mathbf{v}, \mathbf{u} \rangle_{(X^0)'}| \leq |\mathbf{v}|_{(X^0)'}|f\mathbf{u}|_{X^0} \lesssim |\mathbf{v}|_{(X^0)'}(|f|_{L^\infty} + |\nabla f|_{H^2})|\mathbf{u}|_{X^0}.$$

For $n \in \{0, 1\}$, we write

$$|f\mathbf{v}|_{Y^{n+1}} \lesssim |f\mathbf{v}|_{Y^n} + \sum_{|\alpha|=1} |f\partial^\alpha \mathbf{v}|_{Y^n} + |(\partial^\alpha f)\mathbf{v}|_{H^n},$$

where we used (2.2) in Lemma 2.1. Estimate (2.8) follows by finite induction, using (2.4) and again (2.2). For $n > 2$, we use Leibniz rule and the bilinear estimate as above:

$$|(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{Y^0} \lesssim |(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{L^2} \leq |\nabla f|_{H^2}|\mathbf{v}|_{H^{n-1}} + |\nabla f|_{H^{n-1}}|\mathbf{v}|_{H^2}.$$

Estimate (2.8) follows from using again the embedding (2.2). The proof is complete. \square

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Let us now exhibit *elliptic estimates* concerning the operator \mathfrak{T} , defined in (1.2),(1.6).

Lemma 2.3. *Let $b \in \dot{H}^3(\mathbb{R}^d)$ and $h \in \dot{H}^2(\mathbb{R}^d)$ be such that (1.7) holds. Then $\mathfrak{T}[h, \beta b] \in \mathcal{L}(X^0; (X^0)')$ and is symmetric:*

$$\forall \mathbf{u}_1, \mathbf{u}_2 \in X^0, \quad \langle \mathfrak{T}[h, \beta b] \mathbf{u}_1, \mathbf{u}_2 \rangle_{(X^0)'} = \langle \mathfrak{T}[h, \beta b] \mathbf{u}_2, \mathbf{u}_1 \rangle_{(X^0)'}$$

Moreover, one has

$$\forall \mathbf{u}_1, \mathbf{u}_2 \in X^0, \quad |\langle \mathfrak{T}[h, \beta b] \mathbf{u}_1, \mathbf{u}_2 \rangle_{(X^0)'}| \leq C(\mu, h^*, \beta |\nabla b|_{L^\infty}) |\mathbf{u}_1|_{X^0} |\mathbf{u}_2|_{X^0},$$

$$\forall \mathbf{u} \in X^0, \quad |\mathbf{u}|_{X^0}^2 \leq C(h_\star^{-1}) \langle \mathfrak{T}[h, \beta b] \mathbf{u}, \mathbf{u} \rangle_{(X^0)'}$$

In particular, $\mathfrak{T}[h, \beta b] : X^0 \rightarrow (X^0)'$ is a topological isomorphism and

$$\forall \mathbf{v} \in (X^0)', \quad |\mathfrak{T}[h, \beta b]^{-1} \mathbf{v}|_{X^0} \leq C(h_\star^{-1}) |\mathbf{v}|_{(X^0)'}$$

Proof. We establish the estimates for $\mathbf{u}, \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{S}(\mathbb{R}^d)^d$ so that all the terms are well-defined, and the $((X^0)' - X^0)$ duality product coincides with the L^2 inner product. The result for less regular functions is obtained by density of $\mathcal{S}(\mathbb{R}^d)^d$ in X^0 and continuous linear extension.

By definition of \mathfrak{T} in (1.6) and after integration by parts, one has

$$\begin{aligned} (\mathfrak{T}[h, \beta b] \mathbf{u}_1, \mathbf{u}_2)_{L^2} &= \int_{\mathbb{R}^d} h \mathbf{u}_1 \cdot \mathbf{u}_2 + \frac{\mu}{3} h^3 (\nabla \cdot \mathbf{u}_1) (\nabla \cdot \mathbf{u}_2) \\ &- \frac{\mu}{2} h^2 ((\nabla \cdot \mathbf{u}_2) (\beta \nabla b \cdot \mathbf{u}_1) + (\beta \nabla b \cdot \mathbf{u}_2) (\nabla \cdot \mathbf{u}_1)) + \mu h (\beta \nabla b \cdot \mathbf{u}_1) (\beta \nabla b \cdot \mathbf{u}_2), \end{aligned}$$

from which the symmetry is evident. The first estimate of the Lemma follows by Cauchy-Schwarz inequality. The second one is obvious when rewriting

$$(\mathfrak{T}[h, \beta b] \mathbf{u}, \mathbf{u})_{L^2} = \int_{\mathbb{R}^d} h |\mathbf{u}|^2 + \frac{\mu}{12} h^3 |\nabla \cdot \mathbf{u}|^2 + \frac{\mu}{4} h |h \nabla \cdot \mathbf{u} - 2\beta \nabla b \cdot \mathbf{u}|^2.$$

This shows that $\mathfrak{T}[h, \beta b] : X^0 \rightarrow (X^0)'$ is continuous and coercive, so that the operator version of Lax-Milgram theorem ensures that $\mathfrak{T}[h, \beta b]$ is an isomorphism. The continuity of the inverse follows from the coercivity of $\mathfrak{T}[h, \beta b]$:

$$|\mathbf{u}|_{X^0}^2 \leq C(h_\star^{-1}) |\langle \mathfrak{T}[h, \beta b] \mathbf{u}, \mathbf{u} \rangle_{(X^0)'}| \leq C(h_\star^{-1}) |\mathfrak{T}[h, \beta b] \mathbf{u}|_{(X^0)'} |\mathbf{u}|_{X^0},$$

and setting $\mathbf{u} = \mathfrak{T}[h, \beta b]^{-1} \mathbf{v}$ above. \square

Lemma 2.4. *Let $b \in \dot{H}^4(\mathbb{R}^d)$ and $h \in \dot{H}^3(\mathbb{R}^d)$ be such that (1.7) holds; and let $\mathbf{v} \in Y^n$. Then $\mathfrak{T}[h, \beta b]^{-1}\mathbf{v} \in X^n$ and*

$$|\mathfrak{T}[h, \beta b]^{-1}\mathbf{v}|_{X^n} \leq \mathbf{C} \left(|\mathbf{v}|_{Y^n} + \langle |\nabla h|_{H^{n-1}} |\mathbf{v}|_{Y^2} \rangle_{n>2} \right)$$

with $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, |\nabla h|_{H^2}, \beta |\nabla b|_{H^3})$.

Proof. Let $\mathbf{v} \in \mathcal{S}(\mathbb{R}^d)^d$, and denote for simplicity $\mathfrak{T} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]$. One has, for $\Lambda^n = (\text{Id} - \Delta)^{n/2}$,

$$[\Lambda^n, \mathfrak{T}^{-1}]\mathbf{v} = -\mathfrak{T}^{-1}[\Lambda^n, \mathfrak{T}]\mathfrak{T}^{-1}\mathbf{v}.$$

By definition of \mathfrak{T} and since Λ^n commutes with space differentiation, we have for any $\mathbf{u}, \mathbf{w} \in \mathcal{S}(\mathbb{R}^d)^d$,

$$\begin{aligned} |([\Lambda^n, \mathfrak{T}]\mathbf{u}, \mathbf{w})_{L^2}| &= ([\Lambda^n, h]\mathbf{u}, \mathbf{w})_{L^2} + \frac{\mu}{3}([\Lambda^n, h^3]\nabla \cdot \mathbf{u}, \nabla \cdot \mathbf{w})_{L^2} \\ &\quad - \frac{\mu}{2}([\Lambda^n, h^2(\beta \nabla b)]\mathbf{u}, \nabla \cdot \mathbf{w})_{L^2} \\ &\quad - \frac{\mu}{2}([\Lambda^n, h^2(\beta \nabla b)]\nabla \cdot \mathbf{u}, \mathbf{w})_{L^2} \\ &\quad + \mu([\Lambda^n, h(\beta \nabla b)(\beta \nabla b)]\mathbf{u}, \mathbf{w})_{L^2}. \end{aligned}$$

Commutator estimates (see *e.g.* [32, Corollary B.9]) and product estimate (2.5) yield

$$|([\Lambda^n, \mathfrak{T}]\mathbf{u}, \mathbf{w})_{L^2}| \leq \mathbf{C} \left(|\Lambda^{n-1}\mathbf{u}|_{X^0} + \langle |\nabla h|_{H^{n-1}} |\Lambda^2\mathbf{u}|_{X^0} \rangle_{n>2} \right) |\mathbf{w}|_{X^0}.$$

By density and continuity arguments, we infer that for any $\mathbf{u} \in X^{n-1}$, $[\Lambda^n, \mathfrak{T}]\mathbf{u} \in (X^0)'$ and

$$|[\Lambda^n, \mathfrak{T}]\mathbf{u}|_{(X^0)'} \leq \mathbf{C} \left(|\mathbf{u}|_{X^{n-1}} + \langle |\nabla h|_{H^{n-1}} |\mathbf{u}|_{X^2} \rangle_{n>2} \right).$$

Combining the above and by Lemma 2.3, we find

$$\begin{aligned} |\mathfrak{T}^{-1}\mathbf{v}|_{X^n} &= |\mathfrak{T}^{-1}\Lambda^n\mathbf{v} - \mathfrak{T}^{-1}[\Lambda^n, \mathfrak{T}]\mathfrak{T}^{-1}\mathbf{v}|_{X^0} \\ &\leq C(h_\star^{-1})|\Lambda^n\mathbf{v} - [\Lambda^n, \mathfrak{T}]\mathfrak{T}^{-1}\mathbf{v}|_{(X^0)'} \\ &\leq \mathbf{C} \left(|\mathbf{v}|_{Y^n} + |\mathfrak{T}^{-1}\mathbf{v}|_{X^{n-1}} + \langle |\nabla h|_{H^{n-1}} |\mathfrak{T}^{-1}\mathbf{v}|_{X^2} \rangle_{n>2} \right). \end{aligned}$$

The result follows by induction on n , and by density of $\mathcal{S}(\mathbb{R}^d)^d$ in Y^n . \square

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

We now exhibit some expansions of the operators at stake in system (1.3), which are crucial to extract its quasilinear formulation, displayed in Proposition 3.1 below.

Lemma 2.5. *For any $f, g \in \dot{H}^{3\nu n + |\alpha| - 1}(\mathbb{R}^d)$ with $n \in \mathbb{N}$ and $|\alpha| \geq 1$, one has*

$$|\partial^\alpha(fg) - f\partial^\alpha g|_{H^n} \lesssim |\nabla f|_{H^{2\nu n + |\alpha| - 1}} |g|_{H^{n + |\alpha| - 1}}.$$

Proof. Using Leibniz rule, we only have to estimate

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \quad \text{for } |\beta| + |\gamma| \leq n + |\alpha| \text{ and } |\beta| \geq 1.$$

If $|\beta| = 1$, then $|\gamma| \leq n + |\alpha| - 1$, and we find by continuous Sobolev embedding $H^2 \subset L^\infty$

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \lesssim |\partial^\beta f|_{H^2} |\partial^\gamma g|_{L^2}.$$

If $|\beta| = 2$, then $|\gamma| \leq n + |\alpha| - 2$, and standard product estimate (see e.g. [32, Prop. B.2]) yields

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \lesssim |\partial^\beta f|_{H^1} |\partial^\gamma g|_{H^1}.$$

If $|\beta| > 2$, then $|\gamma| \leq n + |\alpha| - 3$, and we find by continuous Sobolev embedding $H^2 \subset L^\infty$

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \lesssim |\partial^\beta f|_{L^2} |\partial^\gamma g|_{H^2}.$$

This concludes the proof. □

Lemma 2.6. *For any $f \in \dot{H}^{4\nu n + |\alpha| - 1}(\mathbb{R}^d)$ and $\mathbf{v} \in Y^{n + |\alpha| - 1}$ with $n \in \mathbb{N}$ and $|\alpha| \geq 1$, one has*

$$|\partial^\alpha(f\mathbf{v}) - f\partial^\alpha \mathbf{v}|_{Y^n} \lesssim |\nabla f|_{H^{3\nu n + |\alpha| - 1}} |\mathbf{v}|_{Y^{n + |\alpha| - 1}}.$$

Proof. Using Leibniz rule, we only have to estimate

$$|(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{Y^0} \quad \text{for } |\beta| + |\gamma| \leq n + |\alpha| \text{ and } |\beta| \geq 1.$$

If $|\beta| = 1$, then $|\gamma| \leq n + |\alpha| - 1$, and we use the product estimate (2.8) in Lemma 2.2 to deduce

$$|(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{Y^0} \lesssim |\partial^\beta f|_{H^3} |\partial^\gamma \mathbf{v}|_{Y^0}.$$

For $|\beta| = 2$, $|\beta| = 3$ and $|\beta| > 3$, we proceed as in the proof of Lemma 2.5, and using (2.2) in Lemma 2.1: by the continuous embedding $L^2 \subset Y^0$ we fall into the L^2 setting, and the continuous embedding $Y^{n + |\alpha| - 1} \subset H^{n + |\alpha| - 2}$ allows to control $|\partial^\gamma \mathbf{v}|_{H^{|\beta| - 2}} \leq |\partial^\gamma \mathbf{v}|_{H^{n + |\alpha| - 2}}$. □

Lemma 2.7. *For any $f, g \in \dot{H}^{3\vee n+|\alpha|-1}(\mathbb{R}^d)$ with $n \in \{0, 1\}$ and $|\alpha| \geq 1$, one has*

$$|\partial^\alpha(fg) - g\partial^\alpha f - f\partial^\alpha g|_{H^n} \lesssim |\nabla f|_{H^2} |\nabla g|_{H^{n+|\alpha|-2}} + |\nabla f|_{H^{n+|\alpha|-2}} |\nabla g|_{H^2}.$$

Proof. The result for $n = 0$ follows from Leibniz rule:

$$\partial^\alpha(fg) - g\partial^\alpha f - f\partial^\alpha g = \sum_{\substack{\beta+\gamma=\alpha \\ |\beta|\geq 1, |\gamma|\geq 1}} \binom{\alpha}{\beta} (\partial^\beta f)(\partial^\gamma g),$$

and we use the standard bilinear estimate [52, Prop. 3.6] to estimate

$$|(\partial^\beta f)(\partial^\gamma g)|_{L^2} \lesssim |\nabla f|_{L^\infty} |\nabla g|_{H^{|\beta|+|\gamma|-2}} + |\nabla f|_{H^{|\beta|+|\gamma|-2}} |\nabla g|_{L^\infty},$$

together with the continuous embedding $H^2 \subset L^\infty$. The case $n = 1$ is obtained in the same way after differentiating the above identity. \square

Lemma 2.8. *For any $f \in \dot{H}^{4\vee n+|\alpha|-1}(\mathbb{R}^d)$, $\mathbf{v} \in Y^{4\vee n+|\alpha|-1}$ with $n \in \{0, 1\}$ and $|\alpha| \geq 1$, one has*

$$|\partial^\alpha(f\mathbf{v}) - \mathbf{v}\partial^\alpha f - f\partial^\alpha \mathbf{v}|_{Y^n} \lesssim |\nabla f|_{H^3} |\mathbf{v}|_{Y^{n+|\alpha|-1}} + |\nabla f|_{H^{n+|\alpha|-2}} |\mathbf{v}|_{Y^4}.$$

Proof. If $\alpha = 1$, then the result is obvious. For $|\alpha| \geq 2$, we use Leibniz rule as above. When $|\beta| = 1$, then by (2.8) in Lemma 2.2,

$$|(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{Y^0} \lesssim (|\partial^\beta f|_{L^\infty} + |\partial^\beta \nabla f|_{H^2}) |\partial^\gamma \mathbf{v}|_{Y^0} \lesssim |\nabla f|_{H^3} |\mathbf{v}|_{Y^{|\alpha|-1}}.$$

When $|\beta| \geq 2$, we use the standard bilinear estimate as above:

$$|(\partial^\beta f)(\partial^\gamma \mathbf{v})|_{L^2} \lesssim |\Lambda \nabla f|_{L^\infty} |\Lambda \mathbf{v}|_{H^{|\beta|+|\gamma|-3}} + |\Lambda \nabla f|_{H^{|\beta|+|\gamma|-3}} |\Lambda \mathbf{v}|_{L^\infty}.$$

This yields the desired estimate for $n = 0$, by the continuous embedding $H^2 \subset L^\infty$ and (2.2) in Lemma 2.1. The case $n = 1$ is obtained similarly. \square

In the following Lemmata, similar expansions are carried on the operators \mathfrak{T} and \mathfrak{T}^{-1} , in particular exhibiting the *shape derivatives* of these operators.

Lemma 2.9. *Let $|\alpha| \geq 1$, $n \in \{0, 1\}$. Let $\zeta \in H^{3\vee n+|\alpha|-1}(\mathbb{R}^d)$, $b \in \dot{H}^{4\vee n+|\alpha|+1}(\mathbb{R}^d)$ be such that (1.7) holds, and $\mathbf{u} \in X^{3\vee n+|\alpha|-1}$. Define*

$$d_h \mathfrak{T}[h, \beta b](f, \mathbf{u}) \stackrel{\text{def}}{=} f\mathbf{u} - \mu \nabla(h^2 f \nabla \cdot \mathbf{u}) + \mu \nabla(fh(\beta \nabla b) \cdot \mathbf{u}) - \mu fh(\beta \nabla b) \nabla \cdot \mathbf{u}. \quad (2.9)$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Then

$$\begin{aligned} & |\partial^\alpha(\mathfrak{T}[h, \beta b]\mathbf{u}) - \mathfrak{T}[h, \beta b]\partial^\alpha\mathbf{u} - \epsilon d_h \mathfrak{T}[h, \beta b](\partial^\alpha\zeta, \mathbf{u})|_{Y^n} \\ & \leq C(\mu, h^*)F(|\nabla b|_{H^{3\nu n+|\alpha|}}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{X^3})\left(|\nabla\zeta|_{H^{n+|\alpha|-2}} + |\mathbf{u}|_{X^{n+|\alpha|-1}}\right). \end{aligned}$$

Proof. Assume first that $\mathbf{u} \in \mathcal{S}(\mathbb{R}^d)^d$, and denote

$$\mathbf{r} \stackrel{\text{def}}{=} \partial^\alpha(\mathfrak{T}[h, \beta b]\mathbf{u}) - \mathfrak{T}[h, \beta b]\partial^\alpha\mathbf{u} - \epsilon d_h \mathfrak{T}[h, \beta b](\partial^\alpha\zeta, \mathbf{u}).$$

By definition of \mathfrak{T} in (1.6), integration by parts and since Λ^n commutes with spatial differentiation, one has for any $\mathbf{w} \in X^0$,

$$\begin{aligned} \langle \Lambda^n \mathbf{r}, \mathbf{w} \rangle_{(X^0)'} &= (\Lambda^n \{ \partial^\alpha(h\mathbf{u}) - h\partial^\alpha\mathbf{u} - \epsilon\partial^\alpha\zeta\mathbf{u} \}, \mathbf{w})_{L^2} \\ &+ \mu \frac{1}{3} (\Lambda^n \{ \partial^\alpha(h^3\nabla \cdot \mathbf{u}) - h^3\partial^\alpha\nabla \cdot \mathbf{u} - 3\epsilon h^2(\partial^\alpha\zeta)\nabla \cdot \mathbf{u} \}, \nabla \cdot \mathbf{w})_{L^2} \\ &- \mu \frac{1}{2} (\Lambda^n \{ \partial^\alpha(h^2(\beta\nabla b) \cdot \mathbf{u}) - h^2(\beta\nabla b) \cdot \partial^\alpha\mathbf{u} - 2\epsilon h(\partial^\alpha\zeta)(\beta\nabla b) \cdot \mathbf{u} \}, \nabla \cdot \mathbf{w})_{L^2} \\ &- \mu \frac{1}{2} (\Lambda^n \{ \partial^\alpha(h^2(\beta\nabla b)\nabla \cdot \mathbf{u}) - h^2(\beta\nabla b)\nabla \cdot \partial^\alpha\mathbf{u} - 2\epsilon h(\partial^\alpha\zeta)(\beta\nabla b)\nabla \cdot \mathbf{u} \}, \mathbf{w})_{L^2} \\ &+ \mu (\Lambda^n \{ \partial^\alpha(h\beta^2(\nabla b \cdot \mathbf{u})\nabla b) - h\beta^2(\nabla b \cdot \partial^\alpha\mathbf{u})\nabla b - \epsilon(\partial^\alpha\zeta)\beta^2(\nabla b \cdot \mathbf{u})\nabla b \}, \mathbf{w})_{L^2} \\ &\stackrel{\text{def}}{=} (\Lambda^n \mathbf{r}^1, \mathbf{w})_{L^2} + \sqrt{\mu} (\Lambda^n \mathbf{r}^2, \nabla \cdot \mathbf{w})_{L^2} \end{aligned}$$

The two contributions are estimated using Lemmata 2.5 and 2.7, product estimates (2.4) and (2.5) and the continuous embedding $H^2 \subset L^\infty$. For instance, by Lemma 2.7, one has

$$|\partial^\alpha(\zeta\mathbf{u}) - \zeta\partial^\alpha\mathbf{u} - \partial^\alpha\zeta\mathbf{u}|_{H^n} \lesssim |\nabla\zeta|_{H^2}|\mathbf{u}|_{H^{n+|\alpha|-1}} + |\nabla\zeta|_{H^{n+|\alpha|-2}}|\mathbf{u}|_{H^3}$$

and by Lemma 2.5,

$$|\partial^\alpha(b\mathbf{u}) - b\partial^\alpha\mathbf{u}|_{H^n} \lesssim |\nabla b|_{H^{2\nu n+|\alpha|-1}}|\mathbf{u}|_{H^{n+|\alpha|-1}},$$

and (2.1) in Lemma 2.1 allows to complete the estimate of the first contribution. The other terms are treated similarly, using first Lemma 2.5 to commute the differentiation operator with bottom contributions, product estimates (2.4) and (2.5) to estimate the commutator, and then Lemma 2.7 to deal with surface contributions.

Altogether, one finds that $|\mathbf{r}^1|_{H^n}$ and $|\mathbf{r}^2|_{H^n}$ are estimated as in the statement. By Cauchy-Schwarz inequality and duality, we deduce that $\mathbf{r} \in Y^n$ and $|\mathbf{r}|_{Y^n}$ satisfies the same estimate. The result for $\mathbf{u} \in X^{3\nu n+|\alpha|-1}$ follows by density and continuous linear extension. \square

Lemma 2.10. *Let $|\alpha| \geq 1$, $n \in \{0, 1\}$. Let $b \in \dot{H}^{4\vee n + |\alpha| + 1}(\mathbb{R}^d)$ and $\zeta \in H^{3\vee n + |\alpha| - 1}(\mathbb{R}^d)$ be such that (1.7) holds, and $\mathbf{v} \in Y^{3\vee n + |\alpha| - 1}$. Denote $\mathfrak{T} = \mathfrak{T}[h, \beta b]$ and $d_h \mathfrak{T} = d_h \mathfrak{T}[h, \beta b]$ defined in (2.9). Then one has*

$$\begin{aligned} & |\partial^\alpha(\mathfrak{T}^{-1}\mathbf{v}) - \mathfrak{T}^{-1}\partial^\alpha\mathbf{v} + \epsilon\mathfrak{T}^{-1}\{d_h\mathfrak{T}(\partial^\alpha\zeta, \mathfrak{T}^{-1}\mathbf{v})\}|_{X^n} \\ & \leq C(\mu, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^{3\vee n + |\alpha|}}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{X^3})\left(|\nabla\zeta|_{H^{n+|\alpha|-2}} + |\mathbf{v}|_{Y^{n+|\alpha|-1}}\right). \end{aligned}$$

Proof. Let us denote $\mathbf{r} = \partial^\alpha(\mathfrak{T}^{-1}\mathbf{v}) - \mathfrak{T}^{-1}\partial^\alpha\mathbf{v} + \epsilon\mathfrak{T}^{-1}\{d_h\mathfrak{T}(\partial^\alpha\zeta, \mathfrak{T}^{-1}\mathbf{v})\}$ and $\mathbf{u} = \mathfrak{T}^{-1}\mathbf{v}$. One has

$$\begin{aligned} \mathbf{r} &= \mathfrak{T}^{-1}\mathfrak{T}\partial^\alpha(\mathfrak{T}^{-1}\mathbf{v}) - \mathfrak{T}^{-1}\partial^\alpha(\mathfrak{T}\mathfrak{T}^{-1}\mathbf{v}) + \epsilon\mathfrak{T}^{-1}\{d_h\mathfrak{T}(\partial^\alpha\zeta, \mathfrak{T}^{-1}\mathbf{v})\} \\ &= -\mathfrak{T}^{-1}\left\{\partial^\alpha(\mathfrak{T}\mathbf{u}) - \mathfrak{T}\partial^\alpha\mathbf{u} - \epsilon d_h\mathfrak{T}(\partial^\alpha\zeta, \mathbf{u})\right\} \stackrel{\text{def}}{=} -\mathfrak{T}^{-1}\tilde{\mathbf{r}}. \end{aligned}$$

By Lemma 2.4 and Lemma 2.9, one has

$$\begin{aligned} |\mathbf{r}|_{X^n} &\leq C(\mu, h_\star^{-1}, h^\star, |\nabla h|_{H^2}, \beta|\nabla b|_{H^3})|\tilde{\mathbf{r}}|_{Y^n} \\ &\leq C(\mu, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^{3\vee n + |\alpha|}}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{X^3}) \\ &\quad \times \left(|\nabla\zeta|_{H^{n+|\alpha|-2}} + |\mathbf{u}|_{X^{n+|\alpha|-1}}\right) \end{aligned}$$

and for $m = 3$ or $m = n + |\alpha| - 1$,

$$|\mathbf{u}|_{X^m} \leq C(\mu, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^2})\left(|\mathbf{v}|_{Y^m} + \langle |\nabla h|_{H^{m-1}}|\mathbf{v}|_{Y^2} \rangle_{m>2}\right).$$

This concludes the proof. \square

We conclude this section with the following result which is essential for estimating the difference between two approximate solutions (in later Proposition 5.2 for instance).

Lemma 2.11. *Let $n \in \mathbb{N}$, $b \in \dot{H}^{4\vee n + 1}(\mathbb{R}^d)$ and $\zeta, \tilde{\zeta} \in H^{3\vee n}(\mathbb{R}^d)$ be such that (1.7) holds, and $\mathbf{v} \in Y^{2\vee n}$. Then, denoting $h = 1 + \epsilon\zeta - \beta b$ and $\tilde{h} = 1 + \epsilon\tilde{\zeta} - \beta b$, one has*

$$|\mathfrak{T}[h, \beta b]^{-1}\mathbf{v} - \mathfrak{T}[\tilde{h}, \beta b]^{-1}\mathbf{v}|_{X^n} \leq \mathbf{C}|\mathbf{v}|_{Y^{2\vee n}}|\zeta - \tilde{\zeta}|_{H^n}$$

with $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, |\nabla b|_{H^{3\vee n}}, |\nabla h|_{H^{2\vee n-1}}, |\nabla\tilde{h}|_{H^{2\vee n-1}})$.

Proof. We rewrite

$$\mathfrak{T}[h, \beta b]^{-1}\mathbf{v} - \mathfrak{T}[\tilde{h}, \beta b]^{-1}\mathbf{v} = \mathfrak{T}[\tilde{h}, \beta b]^{-1}(\mathfrak{T}[\tilde{h}, \beta b] - \mathfrak{T}[h, \beta b])\mathfrak{T}[h, \beta b]^{-1}\mathbf{v}.$$

By Lemma 2.4, one has

$$|\mathfrak{T}[h, \beta b]^{-1} \mathbf{v}|_{X^n} \leq C(\mu, h_\star^{-1}, h^\star, |\nabla h|_{H^{2\nu n-1}}, \beta |\nabla b|_{H^3}) |\mathbf{v}|_{Y^n}.$$

Using the definition (1.6), one has

$$\begin{aligned} \mathfrak{T}[\tilde{h}, \beta b] \mathbf{u} - \mathfrak{T}[h, \beta b] \mathbf{u} &= \epsilon(\zeta - \tilde{\zeta}) \mathbf{u} - \frac{\mu}{3} \nabla((h^3 - \tilde{h}^3) \nabla \cdot \mathbf{u}) \\ &\quad + \frac{\mu}{2} \nabla((h^2 - \tilde{h}^2)(\beta \nabla b \cdot \mathbf{u})) - \frac{\mu}{2} (h - \tilde{h})(\beta \nabla b) \nabla \cdot \mathbf{u}. \end{aligned}$$

By (2.2) and (2.1) in Lemma 2.1 and (2.4) in Lemma 2.2, one has immediately

$$|\epsilon(\zeta - \tilde{\zeta}) \mathbf{u}|_{Y^n} \lesssim \epsilon |\mathbf{u}|_{X^{2\nu n}} |\zeta - \tilde{\zeta}|_{H^n}.$$

Using now (2.3) in Lemma 2.1, we find

$$\begin{aligned} \mu |\nabla((h^3 - \tilde{h}^3) \nabla \cdot \mathbf{u})|_{Y^n} &\leq \sqrt{\mu} |(h^3 - \tilde{h}^3) \nabla \cdot \mathbf{u}|_{H^n} \leq |\mathbf{u}|_{X^{2\nu n}} |h^3 - \tilde{h}^3|_{H^n} \\ &\leq \epsilon C(|\nabla h|_{H^{2\nu n-1}}, |\nabla \tilde{h}|_{H^{2\nu n-1}}) |\mathbf{u}|_{X^{2\nu n}} |\zeta - \tilde{\zeta}|_{H^n}. \end{aligned}$$

The other terms are treated similarly, and we find

$$|\mathfrak{T}[\tilde{h}, \beta b] \mathbf{u} - \mathfrak{T}[h, \beta b] \mathbf{u}|_{Y^n} \leq \mathbf{C} |\mathbf{u}|_{X^{2\nu n}} |\zeta - \tilde{\zeta}|_{H^n}.$$

The proof is completed when collecting the above and applying once again Lemma 2.4. \square

3. The quasilinear system

The result below is the key ingredient of our proof. We extract the quasilinear structure of system (1.3) in terms of “good unknowns”, from which energy estimates will be deduced in subsequent Section 4. The strategy consists in differentiating system (1.3) several times, and estimate lower order contributions thanks to the formulas obtained in Section 2. However, in order not to break the structure of the first equation, we are led (as for the water waves system [27, 32]) to introduce an appropriate velocity variable which is a combination of the original variables and their derivatives.

Proposition 3.1. *Let α be a non-trivial multi-index and $\zeta \in H^{4\nu|\alpha|}(\mathbb{R}^d)$, $b \in \dot{H}^{5\nu|\alpha|+2}(\mathbb{R}^d)$ be such that (1.7) holds, and $\mathbf{v} \in Y^{4\nu|\alpha|}$, satisfying (1.3). Denote $w \stackrel{\text{def}}{=} -h \nabla \cdot \mathbf{u} + \beta \nabla b \cdot \mathbf{u}$ and*

$$\zeta_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta \quad ; \quad \mathbf{v}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \mathbf{v} - \mu \epsilon \nabla(w \partial^\alpha \zeta). \quad (3.1)$$

Then $\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}$ satisfy

$$\begin{cases} \partial_t \zeta_{(\alpha)} + \epsilon \nabla \cdot (\mathbf{u} \zeta_{(\alpha)}) + \nabla \cdot (h \mathbf{u}_{(\alpha)}) = r_{(\alpha)}, \\ (\partial_t + \epsilon \mathbf{u}^\perp \operatorname{curl}) \mathbf{v}_{(\alpha)} + \nabla \zeta_{(\alpha)} + \epsilon \nabla (\mathbf{u} \cdot \mathbf{v}_{(\alpha)}) = \mathbf{r}_{(\alpha)}, \end{cases} \quad (3.2)$$

where we denote

$$\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{I}[h, \beta b]^{-1}(h \mathbf{v}) \quad \text{and} \quad \mathbf{u}_{(\alpha)} \stackrel{\text{def}}{=} \mathfrak{I}[h, \beta b]^{-1}(h \mathbf{v}_{(\alpha)}),$$

and $r_{(\alpha)}, \mathbf{r}_{(\alpha)}$ satisfy the estimates

$$|r_{(\alpha)}|_{L^2} + |\mathbf{r}_{(\alpha)}|_{Y^0} \leq \mathbf{F} \left(|\zeta|_{H^{|\alpha|}} + |\mathbf{v}|_{Y^{|\alpha|}} + |\operatorname{curl} \mathbf{v}|_{H^{|\alpha|-1}} \right) \quad (3.3)$$

with $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^{4\nu|\alpha|+1}}, \epsilon |\nabla \zeta|_{H^3}, \epsilon |\mathbf{v}|_{Y^4}, \epsilon |\operatorname{curl} \mathbf{v}|_{H^3})$.

Moreover, if we denote $\tilde{\zeta}, \tilde{\mathbf{v}}$ satisfying the same assumptions and $\tilde{r}_{(\alpha)}, \tilde{\mathbf{r}}_{(\alpha)}$ the corresponding residuals, then one has

$$\begin{aligned} & |r_{(\alpha)} - \tilde{r}_{(\alpha)}|_{L^2} + |\mathbf{r}_{(\alpha)} - \tilde{\mathbf{r}}_{(\alpha)}|_{Y^0} \\ & \leq \tilde{\mathbf{F}} \left(|\zeta - \tilde{\zeta}|_{H^{|\alpha|}} + |\mathbf{v} - \tilde{\mathbf{v}}|_{Y^{|\alpha|}} + |\operatorname{curl} \mathbf{v} - \operatorname{curl} \tilde{\mathbf{v}}|_{H^{|\alpha|-1}} \right) \end{aligned} \quad (3.4)$$

with

$$\begin{aligned} \tilde{\mathbf{F}} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^{4\nu|\alpha|+1}}, \epsilon |\zeta|_{H^{4\nu|\alpha|}}, \epsilon |\mathbf{v}|_{Y^{4\nu|\alpha|}}, \epsilon |\operatorname{curl} \mathbf{v}|_{H^{3\nu|\alpha|-1}}, \\ \epsilon |\tilde{\zeta}|_{H^{4\nu|\alpha|}}, \epsilon |\tilde{\mathbf{v}}|_{Y^{4\nu|\alpha|}}, \epsilon |\operatorname{curl} \tilde{\mathbf{v}}|_{H^{3\nu|\alpha|-1}}). \end{aligned}$$

Remark 3.2. One recovers the quasilinear structure of the water waves system, as exhibited in [32, Sec. 4.2], up to two differences. Firstly, the advection velocity is \mathbf{u} instead of U (using the notation introduced in Appendix B). This was to be expected, comparing the formulation of the Green-Naghdi system (B.11) with the corresponding formulation of the water waves system (B.12).

Secondly, from the comparison of the aforementioned formulations, one would expect (3.2)₂ to exhibit a component of the form

$$\nabla(\mathbf{a}\zeta) \quad \text{with} \quad \mathbf{a} \stackrel{\text{def}}{=} 1 + \mu \epsilon \partial_t w + \mu \epsilon^2 V \cdot \nabla w,$$

with the hyperbolicity condition $\mathbf{a} > 0$ accounting for the Rayleigh-Taylor criterion, $(-\partial_z P)_{z=\epsilon\zeta} > 0$; see the discussion in [32, Sec. 4.3.5]. In our system, we simply set $\mathbf{a} \equiv 1$ since the additional contributions can be discarded, thanks to our energy norm and the μ prefactor, by (2.3) in

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Lemma 2.1. In other words, using the notation of the proof below, one can check that

$$\nabla \zeta_{(\alpha)} \sim_{Y^0} \nabla \left((1 + \mu \epsilon (\partial_t + \epsilon \mathbf{u} \cdot \nabla) w) \zeta_{(\alpha)} \right).$$

Thus the Rayleigh-Taylor criterion for water waves, which is automatically satisfied for small values of the parameter μ , disappears in the Green-Naghdi system (for any values of μ).

Proof. Let us first remark that by (2.8) in Lemma 2.2, one has for any $\mathbf{v} \in Y^n$ and $n \in \mathbb{N}$,

$$|h\mathbf{v}|_{Y^n} \leq C(h^*)(1 + |\nabla h|_{H^2})|\mathbf{v}|_{Y^n} + \langle |\nabla h|_{H^{n-1}}|\mathbf{v}|_{Y^2} \rangle_{n>2}. \quad (3.5)$$

Using Lemma 2.4, it follows $\mathbf{u} = \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}) \in X^n$ and

$$|\mathbf{u}|_{X^n} \leq C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^3}, \epsilon |\nabla \zeta|_{H^2}) (|\mathbf{v}|_{Y^n} + \langle |\nabla h|_{H^{n-1}}|\mathbf{v}|_{Y^2} \rangle_{n>2}). \quad (3.6)$$

We now enter into the detailed calculations. For simplicity, we denote

$$\begin{aligned} a \sim_{L^2} b &\iff a - b = r, \\ a \sim_{Y^0} b &\iff a - b = \mathbf{r} \end{aligned}$$

with $|r|_{L^2}$ and $|\mathbf{r}|_{Y^0}$ satisfying (3.3).

First equation. We start by differentiating α -times the first equation of (1.3):

$$\partial_t \zeta_{(\alpha)} + \partial^\alpha \nabla \cdot (h\mathbf{u}) = 0.$$

By Lemma 2.7 for the surface contribution and Lemma 2.5 for the bottom contribution (with $n = 1$), and using (2.1) in Lemma 2.1 and (3.6), one finds

$$\partial^\alpha \nabla \cdot (h\mathbf{u}) \sim_{L^2} \nabla \cdot (h\partial^\alpha \mathbf{u}) + \epsilon \nabla \cdot (\mathbf{u}\partial^\alpha \zeta).$$

Now, we use (2.5) in Lemma 2.2, (2.1) in Lemma 2.1, Lemma 2.10 and (3.5), so that

$$\nabla \cdot (h\partial^\alpha \mathbf{u}) \sim_{L^2} \nabla \cdot (h\mathfrak{T}[h, \beta b]^{-1} \{ \partial^\alpha (h\mathbf{v}) - \epsilon d_h \mathfrak{T}[h, \beta b](\partial^\alpha \zeta, \mathbf{u}) \}).$$

Using (2.1) and (2.2) in Lemma 2.1, Lemma 2.4, and proceeding as above but with Lemma 2.8 for the surface contribution and Lemma 2.6 for the bottom contribution, we find

$$\nabla \cdot (h\mathfrak{T}[h, \beta b]^{-1} \{ \partial^\alpha (h\mathbf{v}) \}) \sim_{L^2} \nabla \cdot (h\mathfrak{T}[h, \beta b]^{-1} \{ h\partial^\alpha \mathbf{v} + \epsilon (\partial^\alpha \zeta) \mathbf{v} \}).$$

In order to deal with the second contribution, we use the identity (see the definition (1.6))

$$\mathbf{v} = \mathbf{u} - \frac{\mu}{3h} \nabla(h^3 \nabla \cdot \mathbf{u}) + \frac{\mu}{2h} \nabla(h^2(\beta \nabla b) \cdot \mathbf{u}) - \frac{\mu}{2} h(\beta \nabla b) \nabla \cdot \mathbf{u} + \mu \beta^2 (\nabla b \cdot \mathbf{u}) \nabla b.$$

By the extra μ prefactor, we can use now (2.3) and (2.2) in Lemma 2.1, Lemmata 2.2 and 2.3 as well as (3.6) to deduce

$$\epsilon \nabla \cdot (h \mathfrak{T}[h, \beta b]^{-1} \{(\partial^\alpha \zeta) \mathbf{v}\}) \sim_{L^2} \epsilon \nabla \cdot (h \mathfrak{T}[h, \beta b]^{-1} \{(\partial^\alpha \zeta) \mathbf{u}\}).$$

Finally, recalling (2.9), and proceeding as above, we find

$$\begin{aligned} & \epsilon \nabla \cdot (h \mathfrak{T}[h, \beta b]^{-1} \{d_h \mathfrak{T}[h, \beta b](\partial^\alpha \zeta, \mathbf{u})\}) \\ & \sim_{L^2} \epsilon \nabla \cdot (h \mathfrak{T}[h, \beta b]^{-1} \{(\partial^\alpha \zeta) \mathbf{u} - \mu h \nabla((\partial^\alpha \zeta)(h \nabla \cdot \mathbf{u} - \beta \nabla b \cdot \mathbf{u}))\}). \end{aligned}$$

Collecting the above information and using the definition of $\mathbf{u}_{(\alpha)}$, we obtain, as desired,

$$\partial^\alpha \partial_t \zeta = -\partial^\alpha \nabla \cdot (h \mathbf{u}) \sim_{L^2} -\nabla \cdot (h \mathbf{u}_{(\alpha)}) - \epsilon \nabla \cdot (\mathbf{u} \zeta_{(\alpha)}). \quad (3.7)$$

Second equation. Now we differentiate the second equation of (1.3):

$$\partial_t \partial^\alpha \mathbf{v} + \partial^\alpha (\nabla \zeta + \epsilon \mathbf{u}^\perp \operatorname{curl} \mathbf{v} + \frac{\epsilon}{2} \nabla(|\mathbf{u}|^2)) = \mu \epsilon \partial^\alpha \nabla (\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]).$$

By (2.2) and (2.1) in Lemma 2.1, Lemma 2.7 with $n = 1$ and (3.6), one has

$$\frac{\epsilon}{2} \partial^\alpha \nabla(|\mathbf{u}|^2) \sim_{Y^0} \epsilon \nabla(\mathbf{u} \cdot \partial^\alpha \mathbf{u}).$$

Recalling the definition (1.4)-(1.5), we use (2.3) in Lemma 2.1, Lemma 2.7 with $n = 0$ and Sobolev embedding $H^2 \subset L^\infty$, product estimates (2.5) and (2.6) in Lemma 2.2 and finally (3.6), to deduce

$$\begin{aligned} & \mu \epsilon \partial^\alpha \nabla (\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]) \sim_{Y^0} \\ & \mu \epsilon \nabla \left(\frac{\mathbf{u}}{3h} \cdot \partial^\alpha \nabla(h^3 \nabla \cdot \mathbf{u}) - \frac{1}{2} \frac{\mathbf{u}}{h} \cdot \partial^\alpha \nabla(h^2(\beta \nabla b \cdot \mathbf{u})) \right). \end{aligned}$$

Recalling once again the identity

$$\mathbf{v} = \mathbf{u} - \frac{\mu}{3h} \nabla(h^3 \nabla \cdot \mathbf{u}) + \frac{\mu}{2h} \nabla(h^2(\beta \nabla b) \cdot \mathbf{u}) - \frac{\mu}{2} h(\beta \nabla b) \nabla \cdot \mathbf{u} + \mu \beta^2 (\nabla b \cdot \mathbf{u}) \nabla b,$$

and proceeding as previously for the remainder terms, we deduce

$$\epsilon \partial^\alpha \nabla \left(\frac{1}{2} |\mathbf{u}|^2 - \mu \mathcal{R}[h, \mathbf{u}] - \mu \mathcal{R}_b[h, \beta b, \mathbf{u}] \right) \sim_{Y^0} \epsilon \nabla(\mathbf{u} \cdot \partial^\alpha \mathbf{v}).$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Now, by (2.2) in Lemma 2.1 and Lemma 2.7 with $n = 0$, Sobolev embedding $H^2 \subset L^\infty$ and (3.6), we find

$$\epsilon \partial^\alpha (\mathbf{u}^\perp \operatorname{curl} \mathbf{v}) \sim_{Y^0} \epsilon \mathbf{u}^\perp (\operatorname{curl} \partial^\alpha \mathbf{v}),$$

and the combination yields

$$\partial_t \partial^\alpha \mathbf{v} \sim_{Y^0} -\partial^\alpha \nabla \zeta - \epsilon \nabla (\mathbf{u} \cdot \partial^\alpha \mathbf{v}) - \epsilon \mathbf{u}^\perp (\operatorname{curl} \partial^\alpha \mathbf{v}).$$

In order to conclude, we consider

$$\partial_t (\mathbf{v}_{(\alpha)} - \partial^\alpha \mathbf{v}) = -\mu \epsilon \partial_t \nabla (w \partial^\alpha \zeta).$$

By continuous Sobolev embedding $H^2 \subset L^\infty$, one has

$$|\partial_t w|_{L^\infty} \leq C(h^*, \beta |\nabla b|_{H^2}) (|\partial_t \mathbf{u}|_{X^2} + |\partial_t \zeta|_{H^2}).$$

Using the identity

$$\partial_t \mathbf{u} = \mathfrak{T}[h, \beta b]^{-1} \partial_t (h \mathbf{v}) - \epsilon \mathfrak{T}[h, \beta b]^{-1} \{d_h \mathfrak{T}[h, \beta b] (\partial_t \zeta, \mathbf{u})\},$$

plugging the expressions of $\partial_t \zeta$, $\partial_t \mathbf{v}$ as given by (1.3), and by Lemmata 2.1 and 2.2, Lemma 2.11 with $n = 2$, as well as (3.6), we deduce

$$\partial_t (\mathbf{v}_{(\alpha)} - \partial^\alpha \mathbf{v}) \sim_{Y^0} -\mu \epsilon \nabla (w \partial_t \partial^\alpha \zeta).$$

Finally, using (3.7) and (2.3) in Lemma 2.1, we find after straightforward manipulations and proceeding as above,

$$\begin{aligned} \mu \epsilon \nabla (w \partial_t \partial^\alpha \zeta) &\sim_{Y^0} -\mu \epsilon^2 \nabla (w \nabla \cdot (\mathbf{u} \zeta_{(\alpha)})) - \mu \epsilon \nabla (w \nabla \cdot (h \mathbf{u}_{(\alpha)})) \\ &\sim_{Y^0} -\mu \epsilon^2 \nabla (\mathbf{u} \cdot \nabla (w \zeta_{(\alpha)})). \end{aligned}$$

Collecting the above information yields

$$\begin{aligned} \partial_t \mathbf{v}_{(\alpha)} &\sim_{Y^0} -\partial^\alpha \nabla \zeta - \epsilon \nabla (\mathbf{u} \cdot \partial^\alpha \mathbf{v}) - \epsilon \mathbf{u}^\perp (\operatorname{curl} \partial^\alpha \mathbf{v}) + \mu \epsilon^2 \nabla (\mathbf{u} \cdot \nabla (w \zeta_{(\alpha)})) \\ &= -\partial^\alpha \nabla \zeta - \epsilon \nabla (\mathbf{u} \cdot \mathbf{v}_{(\alpha)}) - \epsilon \mathbf{u}^\perp (\operatorname{curl} \mathbf{v}_{(\alpha)}). \end{aligned} \tag{3.8}$$

Estimates (3.7) and (3.8) provide the first estimate of the statement, namely (3.3).

The second estimate of the statement is obtained identically. Since all contributions on the remainders involve either products or the operator \mathfrak{T}^{-1} , we can express the difference as a sum of terms of the same form but involving at least once $\zeta - \tilde{\zeta}$ or $\mathbf{v} - \tilde{\mathbf{v}}$, or estimated by Lemma 2.11. For instance, we find the corresponding estimate for (3.5) and (3.6) as follows.

Denote $h = 1 + \epsilon\zeta - \beta b$ and $\tilde{h} = 1 + \epsilon\tilde{\zeta} - \beta b$. By (2.2) in Lemma 2.1 and (2.4) and (2.8) in Lemma 2.2,

$$\begin{aligned} |h\mathbf{v} - \tilde{h}\tilde{\mathbf{v}}|_{Y^n} &\leq |h(\mathbf{v} - \tilde{\mathbf{v}})|_{Y^n} + \epsilon|(\zeta - \tilde{\zeta})\tilde{\mathbf{v}}|_{Y^n} \\ &\leq C(\mu, h^*, |\nabla h|_{H^{2\nu n-1}}, \epsilon|\tilde{\mathbf{v}}|_{Y^{3\nu n}})(|\mathbf{v} - \tilde{\mathbf{v}}|_{Y^n} + |\zeta - \tilde{\zeta}|_{H^n}). \end{aligned}$$

Applying this to $\mathbf{u} = \mathfrak{T}^{-1}[h, \beta b](h\mathbf{v})$, $\tilde{\mathbf{u}} = \mathfrak{T}^{-1}[\tilde{h}, \beta b](\tilde{h}\tilde{\mathbf{v}})$ and by Lemma 2.4 and Lemma 2.11, it follows

$$\begin{aligned} |\mathbf{u} - \tilde{\mathbf{u}}|_{X^n} &\leq (|\mathbf{v} - \tilde{\mathbf{v}}|_{Y^n} + |\zeta - \tilde{\zeta}|_{H^n}) \\ &\quad \times C(\mu, h_\star^{-1}, h^*, |\nabla b|_{H^{3\nu n}}, |\nabla h|_{H^{2\nu n-1}}, |\nabla \tilde{h}|_{H^{2\nu n-1}}, \epsilon|\mathbf{v}|_{Y^{2\nu n}}, \epsilon|\tilde{\mathbf{v}}|_{Y^{3\nu n}}). \end{aligned}$$

It is now a tedious but straightforward task to follow the steps of the proof and check that the estimate (3.4) holds. \square

4. A priori energy estimates

This section is dedicated to *a priori* energy estimates on the quasilinearized system (3.2) (or rather a mollified version; see below), which we make use in the proof of the existence and uniqueness of solutions to system (1.3) in subsequent Section 5. It is convenient to introduce the following notation, for $n \in \mathbb{N}$:

$$\mathcal{E}^n(\zeta, \mathbf{v}) \stackrel{\text{def}}{=} |\zeta|_{H^n}^2 + |\mathbf{v}|_{Y^n}^2.$$

We wish to show that the regularity induced by \mathcal{E}^n is propagated by the flow of the Green-Naghdi system (1.3) provided n is chosen sufficiently large. However, the natural energy of our system is determined by the symmetrizer associated with the quasilinear system (3.2) satisfied by $\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}$. This leads us to define

$$\mathcal{F}^n(\zeta, \mathbf{v}) \stackrel{\text{def}}{=} \sum_{0 \leq |\alpha| \leq n} \mathcal{F}[h, \beta b](\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}).$$

with $\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}$ given by (3.1) and

$$\mathcal{F}[h, \beta b](\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) \stackrel{\text{def}}{=} |\zeta_{(\alpha)}|_{L^2}^2 + \langle \mathbf{v}_{(\alpha)}, h\mathbf{u}_{(\alpha)} \rangle_{(X^0)},$$

where we recall that $\mathbf{u}_{(\alpha)} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}_{(\alpha)})$. By convention, we denote $\zeta_{(0)} \stackrel{\text{def}}{=} \zeta$ and $\mathbf{v}_{(0)} \stackrel{\text{def}}{=} \mathbf{v}$. We denote $\mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) = \mathcal{F}[h, \beta b](\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)})$ in the following for the sake of readability.

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

It is not obvious that controlling $\mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)})$ for $0 \leq |\alpha| \leq n$ allows to control $\mathcal{E}^n(\zeta, \mathbf{v})$ and conversely, and this is what we investigate in the following Lemmata.

Lemma 4.1. *Let $b \in \dot{H}^3(\mathbb{R}^d)$ and $\zeta \in H^3(\mathbb{R}^d)$ be such that (1.7) holds, and let $\underline{\zeta} \in L^2(\mathbb{R}^d)$, $\underline{\mathbf{v}} \in Y^0$. Then $\underline{\mathbf{u}} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\underline{\mathbf{v}}) \in X^0$ is uniquely defined, and one has*

$$\mathbf{C}^{-1} |\underline{\mathbf{u}}|_{X^0} \leq |\underline{\mathbf{v}}|_{Y^0} \leq \mathbf{C} |\underline{\mathbf{u}}|_{X^0},$$

and

$$\mathbf{C}^{-1} \mathcal{F}(\underline{\zeta}, \underline{\mathbf{v}}) \leq \mathcal{E}^0(\underline{\zeta}, \underline{\mathbf{v}}) \leq \mathbf{C} \mathcal{F}(\underline{\zeta}, \underline{\mathbf{v}}),$$

with $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2})$.

Proof. The estimates follow from Lemma 2.3 and (2.5),(2.6) and (2.8) in Lemma 2.2. \square

Lemma 4.2. *Let $n \in \mathbb{N}^\star$ and $b \in \dot{H}^4(\mathbb{R}^d)$, $\zeta \in H^{4\nu n}(\mathbb{R}^d)$ be such that (1.7) holds. Assume $\mathbf{v} \in Y^{4\nu n}$ and define $\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}$ as in (3.1). Then one has*

$$\mathcal{F}^n(\zeta, \mathbf{v}) \leq C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^3}, \epsilon^2 \mathcal{E}^4(\zeta, \mathbf{v})) \mathcal{E}^n(\zeta, \mathbf{v}).$$

Conversely, if $\mathbf{v} \in Y^4$ is such that $\mathbf{v}_{(\alpha)} \in Y^0$ for any $1 \leq |\alpha| \leq n$, then $\mathbf{v} \in Y^n$ and

$$\mathcal{E}^n(\zeta, \mathbf{v}) \leq C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^3}, \epsilon^2 \mathcal{F}^4(\zeta, \mathbf{v})) \mathcal{F}^n(\zeta, \mathbf{v}).$$

Proof. By Lemma 4.1, it suffices to prove the estimates replacing \mathcal{F}^n with

$$\tilde{\mathcal{E}}^n(\zeta, \mathbf{v}) \stackrel{\text{def}}{=} \sum_{0 \leq |\alpha| \leq n} \mathcal{E}^0(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}).$$

Recall definition (3.1): $\zeta_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta$ and $\mathbf{v}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \mathbf{v} - \mu \epsilon \nabla(w \partial^\alpha \zeta)$ with $w \stackrel{\text{def}}{=} -h \nabla \cdot \mathbf{u} + \beta \nabla b \cdot \mathbf{u}$. By Lemma 2.2 and Lemma 2.4, one has for any $m \in \{0, 1, 2\}$,

$$\begin{aligned} \sqrt{\mu} |w|_{H^m} &\leq C(\mu, h^\star, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}) |\mathbf{u}|_{X^m} \\ &\leq C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^3}, \epsilon |\zeta|_{H^3}) |\mathbf{v}|_{Y^m}. \end{aligned}$$

Now, by (2.3) in Lemma 2.1 and continuous Sobolev embedding $H^2 \subset L^\infty$, one has

$$\mu \epsilon |\nabla(w \zeta_{(\alpha)})|_{Y^0} \lesssim \epsilon \sqrt{\mu} |w \zeta_{(\alpha)}|_{L^2} \lesssim \epsilon \sqrt{\mu} |w|_{H^2} |\zeta_{(\alpha)}|_{L^2}.$$

We immediately deduce the first inequality of the statement.

For the converse, we first use that if $|\alpha| \leq 2$, one has

$$\mu\epsilon|\nabla(w\zeta(\alpha))|_{Y^0} \lesssim \epsilon\sqrt{\mu}|w\zeta(\alpha)|_{L^2} \lesssim \epsilon\sqrt{\mu}|\zeta(\alpha)|_{H^2}|w|_{L^2} \leq \sqrt{\mu}|w|_{L^2}|\epsilon\zeta|_{H^4}.$$

This yields for $n \in \{0, 1, 2\}$,

$$|\mathbf{v}|_{Y^n}^2 \leq C(\mu, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^3}, \epsilon|\zeta|_{H^4}) \times \left(\sum_{0 \leq |\alpha| \leq n} |\mathbf{v}(\alpha)|_{Y^0}^2 \right).$$

Then, for $|\alpha| > 2$, we use as previously

$$\mu\epsilon|\nabla(w\zeta(\alpha))|_{Y^0} \lesssim \epsilon\sqrt{\mu}|w\zeta(\alpha)|_{L^2} \lesssim \epsilon\sqrt{\mu}|w|_{H^2}|\zeta(\alpha)|_{L^2}.$$

We deduce that for $n \geq 3$,

$$|\mathbf{v}|_{Y^n}^2 \leq C(\mu, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^3}, \epsilon|\zeta|_{H^4}, \epsilon|\mathbf{v}|_{Y^2}) \times \left(\sum_{0 \leq |\alpha| \leq n} \mathcal{E}^0(\zeta(\alpha), \mathbf{v}(\alpha)) \right),$$

so that the second inequality of the statement follows. \square

In the following, we provide energy estimates for regular solutions of a mollified version of the Green-Naghdi system (1.3), namely

$$\begin{cases} \partial_t \zeta + J^\iota \nabla \cdot (h\mathbf{u}) = r^\iota, \\ \partial_t \mathbf{v} + J^\iota \left(\nabla \zeta + \epsilon \mathbf{u}^\perp \operatorname{curl} \mathbf{v} + \frac{\epsilon}{2} \nabla (|\mathbf{u}|^2) - \mu \epsilon \nabla (\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]) \right) = \mathbf{r}^\iota, \end{cases} \quad (4.1)$$

where $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v})$ as well as its associated quasilinear system (see Proposition 3.1)

$$\begin{cases} \partial_t \zeta(\alpha) + J^\iota \left(\epsilon \nabla \cdot (\mathbf{u}\zeta(\alpha)) + \nabla \cdot (h\mathbf{u}(\alpha)) \right) = r^\iota(\alpha), \\ \partial_t \mathbf{v}(\alpha) + J^\iota \left(\nabla \zeta(\alpha) + \epsilon \mathbf{u}^\perp \operatorname{curl} \mathbf{v}(\alpha) + \epsilon \nabla (\mathbf{u} \cdot \mathbf{v}(\alpha)) \right) = \mathbf{r}^\iota(\alpha), \end{cases} \quad (4.2)$$

where we denote $\mathbf{u}(\alpha) \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}(\alpha))$. Here and thereafter, J^ι is a Friedrichs mollifier, defined as the Fourier multiplier $J^\iota = \varphi(\iota|D|)$, where $\iota \in (0, 1)$ is a parameter and φ is a smooth function taking values in $[0, 1]$, compactly supported and equal to 1 in a neighborhood of the origin. By convention we write $J^0 \equiv \text{Id}$ and $J^\iota(u_1, u_2) = (J^\iota u_1, J^\iota u_2)$. The following properties will be used repeatedly:

Lemma 4.3. *Let $n \leq m \in \mathbb{N}$, $\iota, \iota_1, \iota_2 \in (0, 1)$, and define the Fourier multiplier J^ι as above. Then for any $f \in H^n(\mathbb{R}^d)$ and $\mathbf{v} \in Y^n$, one has $J^\iota f \in H^m(\mathbb{R}^d)$, $J^\iota \mathbf{v} \in Y^m$ and*

$$|f - J^\iota f|_{H^n} + \iota^{-1} |f - J^\iota f|_{H^{n-1}} + |\mathbf{v} - J^\iota \mathbf{v}|_{Y^n} + \iota^{-1} |\mathbf{v} - J^\iota \mathbf{v}|_{Y^{n-1}} \rightarrow 0 \quad (\iota \rightarrow 0). \quad (4.3)$$

The above holds for $f_n \rightarrow f$ in H^n and $\mathbf{v}_n \rightarrow \mathbf{v}$ in Y^n , uniformly with respect to $n \in \mathbb{N}$. Moreover, there exists a constant C , depending only on φ , such that

$$|J^\iota f|_{H^n} \leq |f|_{H^n}, \quad |J^\iota \mathbf{v}|_{Y^n} \leq |\mathbf{v}|_{Y^n}, \quad (4.4)$$

$$|J^\iota f|_{H^m} \leq C \iota^{n-m} |f|_{H^n}, \quad |J^\iota \mathbf{v}|_{Y^m} \leq C \iota^{n-m} |\mathbf{v}|_{Y^n}, \quad (4.5)$$

$$|(J^{\iota_2} - J^{\iota_1})f|_{H^{n-1}} + |(J^{\iota_2} - J^{\iota_1})\mathbf{v}|_{Y^{n-1}} \leq C |\iota^2 - \iota^1| (|f|_{H^n} + |\mathbf{v}|_{Y^n}). \quad (4.6)$$

Moreover, if $f \in H^3(\mathbb{R}^d)$ and $g \in H^{n-1}(\mathbb{R}^d)$, there exists C_n , depending only on φ and n such that

$$|[J^\iota, f]g|_{H^n} \leq C |\nabla f|_{H^{2\nu_{n-1}}} |g|_{H^{n-1}}. \quad (4.7)$$

Proof. The first estimates in H^n are straightforward by Fourier analysis (see [9, Lemma 5]). The estimates in Y^n follow by duality and using that J^ι is symmetric and commutes with spatial derivatives. The last estimate is a consequence of [32, Corollary B.9] and (4.4). \square

Propositions 4.4 and 4.5 below provide *a priori* energy inequalities for sufficiently regular solutions of (4.1) and (4.2). Again, these estimates are uniform with respect to $\iota \in (0, 1)$, and in particular hold as well for $J^\iota = \text{Id}$.

Proposition 4.4. *Let $b \in \dot{H}^4(\mathbb{R}^d)$, $(\zeta, \mathbf{v}) \in C^1([0, T]; H^3(\mathbb{R}^d)^{1+d})$ be such that (1.7) holds and satisfy system (4.1) with residuals $(r^\iota, \mathbf{r}^\iota) \in C^0([0, T]; L^2(\mathbb{R}^d)^{1+d})$. Then for any $t \in [0, T]$,*

$$\frac{d}{dt} \mathcal{F}(\zeta, \mathbf{v}) \leq \mathbf{F} \mathcal{F}(\zeta, \mathbf{v}) + \mathbf{C} (\mathcal{F}(r^\iota, \mathbf{r}^\iota) \mathcal{F}(\zeta, \mathbf{v}))^{1/2}$$

where we denote $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^3}, \epsilon |\partial_t \zeta|_{H^3}, \epsilon |\nabla \zeta|_{H^2}, \epsilon |\mathbf{v}|_{H^3})$ and $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2})$, independent of $\iota \in (0, 1)$.

Proof. The regularity assumptions on the data and the assumption (1.7) are sufficient to ensure that all the terms and calculations (including integration by parts) below are well-defined. We test the first equation of the

system (4.1) against $\zeta \in C^0([0, T]; L^2(\mathbb{R}^d))$, and the second one against $h\mathbf{u} = h\mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}) \in C^0([0, T]; X^0)$. It follows

$$\frac{1}{2} \frac{d}{dt} \mathcal{F}(\zeta, \mathbf{v}) = A^1 + A^{2,\iota} + A^{3,\iota} + A^4$$

with, using the symmetry of the operator $\mathfrak{T}[h, \beta b]^{-1}$,

$$A^1 \stackrel{\text{def}}{=} \frac{1}{2} (\mathbf{v}, [\partial_t, h\mathfrak{T}[h, \beta b]^{-1}h]\mathbf{v})_{L^2}$$

$$A^{2,\iota} \stackrel{\text{def}}{=} -\epsilon (J^\iota(\mathbf{u}^\perp \operatorname{curl} \mathbf{v} + \frac{1}{2} \nabla(|\mathbf{u}|^2) - \mu \nabla(\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}])), h\mathbf{u})_{L^2}$$

$$A^{3,\iota} \stackrel{\text{def}}{=} -(J^\iota \nabla \cdot (h\mathbf{u}), \zeta)_{L^2} - (J^\iota \nabla \zeta, h\mathbf{u})_{L^2}$$

$$A^4 \stackrel{\text{def}}{=} (r^\iota, \zeta)_{L^2} + (\mathbf{r}^\iota, h\mathbf{u})_{L^2}.$$

Since J^ι is symmetric and commutes with differential operators, one has $A^{3,\iota} \equiv 0$ after integrating by parts. The contributions of A^1 and A^4 are treated in details in the proof of Proposition 4.5; see (4.8) and (4.12), below, with Lemma 4.1. As for $A^{2,\iota}$, one obtains rough estimates as follows.

By continuous Sobolev embedding $H^2 \subset L^\infty$ and (2.5) in Lemma 2.2,

$$|\epsilon \nabla \cdot (h\mathbf{u})|_{L^\infty} \lesssim |h \times (\epsilon \mathbf{u})|_{H^3} \leq C(h^*) F(\beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}, \epsilon |\mathbf{u}|_{H^3})$$

and therefore, by (4.4) in Lemma 4.3 and Cauchy-Schwarz inequality,

$$|\epsilon (J^\iota(\frac{1}{2} |\mathbf{u}|^2), \nabla \cdot (h\mathbf{u}))_{L^2}| \leq C(h^*) F(\beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}, \epsilon |\mathbf{u}|_{H^3}) |\mathbf{u}|_{L^2}^2.$$

Similarly, recalling the definition (1.4)-(1.5), one obtains as above

$$\sqrt{\mu} \epsilon |\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]|_{L^2} \leq C(\mu, h_\star^{-1}, h^*) F(\beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}, \epsilon |\mathbf{u}|_{X^3}) |\mathbf{u}|_{X^0}$$

and

$$\begin{aligned} \sqrt{\mu} |\nabla \cdot (h\mathbf{u})|_{L^2} &\leq \sqrt{\mu} |\nabla h|_{L^\infty} |\mathbf{u}|_{L^2} + \sqrt{\mu} |h|_{L^\infty} |\nabla \cdot \mathbf{u}|_{L^2} \\ &\leq C(\mu, h^*, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}) |\mathbf{u}|_{X^0}. \end{aligned}$$

Finally, one has by Cauchy-Schwarz inequality and continuous Sobolev embedding $H^2 \subset L^\infty$

$$|(J^\iota(\mathbf{u}^\perp \operatorname{curl} \mathbf{v}), h\mathbf{u})_{L^2}| \leq C(h^*) |\operatorname{curl} \mathbf{v}|_{H^2} |\mathbf{u}|_{L^2}^2.$$

It follows, by (2.1) in Lemma 2.1,

$$|A^{2,\iota}| \leq C(\mu, h_\star^{-1}, h^*) F(\beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}, \epsilon |\mathbf{u}|_{X^3}, \epsilon |\operatorname{curl} \mathbf{v}|_{H^2}) |\mathbf{u}|_{X^0}^2.$$

We conclude by Lemma 2.4 and (2.8) in Lemma 2.2, and Lemma 4.1. \square

Proposition 4.5. *Let $(\zeta, \mathbf{u}) \in C^0([0, T]; H^4(\mathbb{R}^d)^{1+d}) \cap C^1([0, T]; H^3(\mathbb{R}^d)^{1+d})$ and $b \in \dot{H}^4$ be such that (1.7) holds, and $(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) \in C^0([0, T]; H^1(\mathbb{R}^d)^{1+d})$ satisfying system (4.2) with remainders $(r_{(\alpha)}^\iota, \mathbf{r}_{(\alpha)}^\iota) \in C^0([0, T]; L^2(\mathbb{R}^d)^{1+d})$. Then one has for any $t \in [0, T]$,*

$$\frac{d}{dt} \mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) \leq \mathbf{F} \mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) + \mathbf{C} (\mathcal{F}(r_{(\alpha)}^\iota, \mathbf{r}_{(\alpha)}^\iota) \mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}))^{1/2}$$

where we denote $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^3}, \epsilon |\partial_t \zeta|_{H^3}, \epsilon |\nabla \zeta|_{H^3}, \epsilon |\mathbf{u}|_{H^3})$ and $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2})$, independent of $\iota \in (0, 1)$.

Proof. The regularity assumptions on the data and the assumption (1.7) are sufficient to ensure that all the terms and calculations below are well-defined. We test the first equation of (4.2) against $\zeta_{(\alpha)} \in C^0([0, T]; L^2)$ and the second against $h\mathbf{u}_{(\alpha)} = h\mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}_{(\alpha)}) \in C^0([0, T]; X^0)$. It follows

$$\frac{1}{2} \frac{d}{dt} \mathcal{F}(\zeta_{(\alpha)}, \mathbf{v}_{(\alpha)}) = A_{(\alpha)}^1 + A_{(\alpha)}^{2,\iota} + A_{(\alpha)}^{3,\iota} + A_{(\alpha)}^4$$

with, using the symmetry of the operator $\mathfrak{T}[h, \beta b]^{-1}$,

$$A_{(\alpha)}^1 \stackrel{\text{def}}{=} \frac{1}{2} (\mathbf{v}_{(\alpha)}, [\partial_t, h\mathfrak{T}[h, \beta b]^{-1}h] \mathbf{v}_{(\alpha)})_{L^2}$$

$$A_{(\alpha)}^{2,\iota} \stackrel{\text{def}}{=} -\epsilon (J^\iota \nabla \cdot (\mathbf{u} \zeta_{(\alpha)}), \zeta_{(\alpha)})_{L^2} - \epsilon (J^\iota (\mathbf{u}^\perp \text{curl } \mathbf{v}_{(\alpha)} + \nabla(\mathbf{u} \cdot \mathbf{v}_{(\alpha)})), h\mathbf{u}_{(\alpha)})_{L^2}$$

$$A_{(\alpha)}^{3,\iota} \stackrel{\text{def}}{=} -(J^\iota \nabla \cdot (h\mathbf{u}_{(\alpha)}), \zeta_{(\alpha)})_{L^2} - (J^\iota \nabla \zeta_{(\alpha)}, h\mathbf{u}_{(\alpha)})_{L^2}$$

$$A_{(\alpha)}^4 \stackrel{\text{def}}{=} (r_{(\alpha)}^{1,\iota}, \zeta_{(\alpha)})_{L^2} + (r_{(\alpha)}^{2,\iota}, h\mathbf{u}_{(\alpha)})_{L^2}.$$

We now estimate each contribution in terms of $|\mathbf{v}_{(\alpha)}|_{Y^0}$, $|\mathbf{u}_{(\alpha)}|_{X^0}$, $|\zeta_{(\alpha)}|_{L^2}$. *Estimate on $A_{(\alpha)}^1$.* We use the explicit formula for the commutator

$$\begin{aligned} & \frac{1}{2} (\mathbf{v}_{(\alpha)}, [\partial_t, h\mathfrak{T}[h, \beta b]^{-1}h] \mathbf{v}_{(\alpha)})_{L^2} \\ &= (\mathbf{v}_{(\alpha)}, (\partial_t h) \mathfrak{T}[h, \beta b]^{-1} \{h\mathbf{v}_{(\alpha)}\})_{L^2} + \frac{1}{2} (\mathbf{v}_{(\alpha)}, h[\partial_t, \mathfrak{T}[h, \beta b]^{-1}] h \mathbf{v}_{(\alpha)})_{L^2} \\ &= ((\partial_t h) \mathbf{v}_{(\alpha)}, \mathbf{u}_{(\alpha)})_{L^2} - \frac{1}{2} (\mathbf{u}_{(\alpha)}, [\partial_t, \mathfrak{T}[h, \beta b]] \mathbf{u}_{(\alpha)})_{L^2} \\ &= ((\partial_t h) \mathbf{v}_{(\alpha)}, \mathbf{u}_{(\alpha)})_{L^2} - \frac{1}{2} (\mathbf{u}_{(\alpha)}, (\partial_t h) \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad - \frac{\mu}{2} (\nabla \cdot \mathbf{u}_{(\alpha)}, (h^2 \partial_t h) \nabla \cdot \mathbf{u}_{(\alpha)})_{L^2} + \mu ((h \partial_t h) \nabla \cdot \mathbf{u}_{(\alpha)}, (\beta \nabla b) \cdot \mathbf{u}_{(\alpha)})_{L^2} \end{aligned}$$

where we used the symmetry of $\mathfrak{T}[h, \beta b]^{-1}$ and the definitions (1.6),(1.2). Since $\partial_t h = \epsilon \partial_t \zeta$, we deduce by (2.8) in Lemma 2.2, continuous Sobolev embedding $H^2 \subset L^\infty$ and Cauchy-Schwarz inequality,

$$|A_{(\alpha)}^1| \leq C(\mu, h^*, \beta |\nabla b|_{H^2}) \times (|\epsilon \partial_t \zeta|_{H^3} |\mathbf{u}_{(\alpha)}|_{X^0} |\mathbf{v}_{(\alpha)}|_{Y^0} + |\epsilon \partial_t \zeta|_{H^2} |\mathbf{u}_{(\alpha)}|_{X^0}^2). \quad (4.8)$$

Estimate on $A_{(\alpha)}^{2,\iota}$. First remark that, since J^ι is symmetric and commutes with differential operators, one has after integrating by parts

$$A_{(\alpha)}^{2,\iota,i} \stackrel{\text{def}}{=} -\epsilon (J^\iota \nabla \cdot (\mathbf{u} \zeta_{(\alpha)}), \zeta_{(\alpha)})_{L^2} = \epsilon (\zeta_{(\alpha)}, \mathbf{u} \cdot \nabla J^\iota \zeta_{(\alpha)})_{L^2}$$

and therefore, averaging the left-hand side and the right-hand side,

$$A_{(\alpha)}^{2,\iota,i} \stackrel{\text{def}}{=} -\frac{1}{2} \epsilon (J^\iota (\zeta_{(\alpha)} \nabla \cdot \mathbf{u}), \zeta_{(\alpha)})_{L^2} - \frac{1}{2} \epsilon (\zeta_{(\alpha)}, [J^\iota, \mathbf{u}] \nabla \zeta_{(\alpha)})_{L^2}.$$

By the product and commutator estimates (4.4) and (4.7) in Lemma 4.3, and applying Cauchy-Schwarz inequality and continuous Sobolev embedding $H^2 \subset L^\infty$, we get

$$|A_{(\alpha)}^{2,\iota,i}| \lesssim \epsilon |\mathbf{u}|_{H^3} \times |\zeta_{(\alpha)}|_{L^2}^2, \quad (4.9)$$

uniformly with respect to $\iota \in (0, 1)$.

The second contribution, namely

$$A_{(\alpha)}^{2,\iota,ii} \stackrel{\text{def}}{=} (J^\iota (\mathbf{u}^\perp \text{curl } \mathbf{v}_{(\alpha)} + \nabla (\mathbf{u} \cdot \mathbf{v}_{(\alpha)})), h \mathfrak{T}[h, \beta b]^{-1} (h \mathbf{v}_{(\alpha)}))_{L^2},$$

is by far the most involved (notice that in the case of the water waves system, the corresponding term requires a specific attention as well; see [32, Prop. 3.30]). We prove in Lemma 4.6, below, that

$$|A_{(\alpha)}^{2,\iota,ii}| \leq C(\mu, h_\star^{-1}, h^*) F(\beta |\nabla b|_{H^3}, \epsilon |\nabla \zeta|_{H^3}, \epsilon |\mathbf{u}|_{H^3}) \times |\mathbf{v}_{(\alpha)}|_{Y^0}^2. \quad (4.10)$$

Estimate on $A_{(\alpha)}^{3,\iota}$. Thanks to our choice of the symmetrizer and since J^ι is symmetric and commutes with differential operators, one has after integrating by parts

$$A_{(\alpha)}^{3,\iota} \equiv 0. \quad (4.11)$$

Estimate on $A_{(\alpha)}^4$. By Cauchy-Schwarz inequality and (2.7) in Lemma 2.2,

$$|A_{(\alpha)}^4| \leq C(h^*, \beta |\nabla b|_{H^2}, \epsilon |\nabla \zeta|_{H^2}) \times (|\zeta_{(\alpha)}|_{L^2}^2 + |\mathbf{u}_{(\alpha)}|_{X^0}^2)^{1/2} \mathcal{E}^0(r_{(\alpha)}^1, r_{(\alpha)}^2). \quad (4.12)$$

Estimates (4.8)–(4.12), with Lemma 4.1 yield the desired result. \square

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

We conclude this section with the proof of estimate (4.10), which is essential in the proof of Proposition 4.5.

Lemma 4.6. *Under the assumptions of Proposition 4.5, one has*

$$\epsilon |(J^\iota(\mathbf{u}^\perp \operatorname{curl} \mathbf{v}_{(\alpha)} + \nabla(\mathbf{u} \cdot \mathbf{v}_{(\alpha)})), h\mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}_{(\alpha)}))_{L^2}| \leq \mathbf{F} |\mathbf{v}_{(\alpha)}|_{Y^0}^2,$$

with $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^3}, \epsilon|\mathbf{u}|_{H^3})$, uniformly with respect to $\iota \in (0, 1)$.

Proof. We denote $\mathbf{u}_{(\alpha)} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}_{(\alpha)})$ and use the identity valid for any U and V sufficiently regular two-dimensional vector fields

$$\nabla(U \cdot V) = (U \cdot \nabla)V + (V \cdot \nabla)U - V^\perp \operatorname{curl} U - U^\perp \operatorname{curl} V. \quad (4.13)$$

Thus we have

$$\begin{aligned} B^\iota &\stackrel{\text{def}}{=} (J^\iota(\mathbf{u}^\perp \operatorname{curl} \mathbf{v}_{(\alpha)} + \nabla(\mathbf{u} \cdot \mathbf{v}_{(\alpha)})), h\mathbf{u}_{(\alpha)})_{L^2} \\ &= ((\mathbf{u} \cdot \nabla)\mathbf{v}_{(\alpha)} + (\mathbf{v}_{(\alpha)} \cdot \nabla)\mathbf{u} - \mathbf{v}_{(\alpha)}^\perp \operatorname{curl} \mathbf{u}, J^\iota(h\mathbf{u}_{(\alpha)}))_{L^2} \\ &= -((\mathbf{u} \cdot \nabla h)\mathbf{v}_{(\alpha)}, J^\iota \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + ((\mathbf{u} \cdot \nabla)(h\mathbf{v}_{(\alpha)}) + (h\mathbf{v}_{(\alpha)} \cdot \nabla)\mathbf{u} - (h\mathbf{v}_{(\alpha)})^\perp \operatorname{curl} \mathbf{u}, J^\iota \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + ((\mathbf{u} \cdot \nabla)\mathbf{v}_{(\alpha)} + (\mathbf{v}_{(\alpha)} \cdot \nabla)\mathbf{u} - \mathbf{v}_{(\alpha)}^\perp \operatorname{curl} \mathbf{u}, [J^\iota, h]\mathbf{u}_{(\alpha)})_{L^2}. \end{aligned}$$

The first term in B^ι is controlled by Cauchy-Schwarz inequality, (2.4) and (2.8) in Lemma 2.2 and (4.4) in Lemma 4.3:

$$\begin{aligned} \epsilon |B_0^\iota| &\stackrel{\text{def}}{=} \epsilon |((\mathbf{u} \cdot \nabla h)\mathbf{v}_{(\alpha)}, J^\iota \mathbf{u}_{(\alpha)})_{L^2}| \\ &\leq F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^3}, \epsilon|\mathbf{u}|_{H^3}) |\mathbf{v}_{(\alpha)}|_{Y^0} |\mathbf{u}_{(\alpha)}|_{X^0}. \end{aligned} \quad (4.14)$$

For the second term, we plug the identity (recall the definition of $\mathfrak{T}[h, \beta b]$ in (1.2) and (1.6))

$$\begin{aligned} h\mathbf{v}_{(\alpha)} &= h\mathbf{u}_{(\alpha)} - \frac{\mu}{3} \nabla(h^3 \nabla \cdot \mathbf{u}_{(\alpha)}) + \frac{\mu}{2} \left(\nabla(h^2(\beta \nabla b) \cdot \mathbf{u}_{(\alpha)}) - h^2(\beta \nabla b) \nabla \cdot \mathbf{u}_{(\alpha)} \right) \\ &\quad + \mu h \beta^2 (\nabla b \cdot \mathbf{u}_{(\alpha)}) \nabla b \end{aligned}$$

and consider separately the four contributions.

One has

$$B_1^\iota \stackrel{\text{def}}{=} ((\mathbf{u} \cdot \nabla)(h\mathbf{u}_{(\alpha)}), J^\iota \mathbf{u}_{(\alpha)})_{L^2} + ((h\mathbf{u}_{(\alpha)} \cdot \nabla)\mathbf{u} - (h\mathbf{u}_{(\alpha)})^\perp \operatorname{curl} \mathbf{u}, J^\iota \mathbf{u}_{(\alpha)})_{L^2}.$$

Integrating by parts the advection operator and averaging yields

$$\begin{aligned} ((\mathbf{u} \cdot \nabla)(h\mathbf{u}_{(\alpha)}), J^t \mathbf{u}_{(\alpha)})_{L^2} &= \frac{1}{2}(\mathbf{u}_{(\alpha)}, [J^t, (h\mathbf{u} \cdot \nabla)]\mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + \frac{1}{2}((\mathbf{u} \cdot \nabla h)\mathbf{u}_{(\alpha)}, J^t \mathbf{u}_{(\alpha)})_{L^2} - \frac{1}{2}(\mathbf{u}_{(\alpha)}, (h\nabla \cdot \mathbf{u})J^t \mathbf{u}_{(\alpha)})_{L^2}. \end{aligned}$$

All components are now estimated by Cauchy-Schwarz inequality and continuous Sobolev embedding $H^2 \subset L^\infty$ as well as (4.4),(4.7) in Lemma 4.3:

$$\epsilon|B_1^t| \leq C(h^*)F(\beta|\nabla b|_{H^2}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{u}_{(\alpha)}|_{L^2}^2. \quad (4.15)$$

One has, denoting $V = \nabla(h^3\nabla \cdot \mathbf{u}_{(\alpha)})$,

$$\begin{aligned} B_2^t &\stackrel{\text{def}}{=} -\frac{\mu}{3}((\mathbf{u} \cdot \nabla)V + (V \cdot \nabla)\mathbf{u} - (V^\perp \text{curl } \mathbf{u}, J^t \mathbf{u}_{(\alpha)}))_{L^2} \\ &= \frac{\mu}{3}(\mathbf{u} \cdot V, J^t \nabla \cdot \mathbf{u}_{(\alpha)})_{L^2} \\ &= -\frac{\mu}{6}((h^3\nabla \cdot \mathbf{u} - 3h^2\mathbf{u} \cdot \nabla h)\nabla \cdot \mathbf{u}_{(\alpha)}, J^t \nabla \cdot \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + \frac{\mu}{6}(\nabla \cdot \mathbf{u}_{(\alpha)}, [J^t, h^3\mathbf{u}] \cdot \nabla(\nabla \cdot \mathbf{u}_{(\alpha)}))_{L^2}, \end{aligned}$$

where we used the identity (4.13) (notice that $\text{curl } V = 0$) and integration by parts. We conclude as above

$$\epsilon|B_2^t| \leq \mu C(h^*)F(\beta|\nabla b|_{H^2}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{H^3})|\nabla \cdot \mathbf{u}_{(\alpha)}|_{L^2}^2. \quad (4.16)$$

One has, denoting for readability $F \stackrel{\text{def}}{=} h^2(\beta\nabla b)$ and $V = \nabla(F \cdot \mathbf{u}_{(\alpha)})$,

$$\begin{aligned} B_3^t &\stackrel{\text{def}}{=} \frac{\mu}{2}((\mathbf{u} \cdot \nabla)V + (V \cdot \nabla)\mathbf{u} - (V^\perp \text{curl } \mathbf{u}, J^t \mathbf{u}_{(\alpha)}))_{L^2} \\ &\quad - \frac{\mu}{2}((\mathbf{u} \cdot \nabla)(F\nabla \cdot \mathbf{u}_{(\alpha)}) + ((F\nabla \cdot \mathbf{u}_{(\alpha)}) \cdot \nabla)\mathbf{u} - (F\nabla \cdot \mathbf{u}_{(\alpha)})^\perp \text{curl } \mathbf{u}, J^t \mathbf{u}_{(\alpha)})_{L^2} \\ &= -\frac{\mu}{2}(\mathbf{u} \cdot V, J^t \nabla \cdot \mathbf{u}_{(\alpha)})_{L^2} - \frac{\mu}{2}((\mathbf{u} \cdot \nabla)(F\nabla \cdot \mathbf{u}_{(\alpha)}), J^t \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad - \frac{\mu}{2}(((F\nabla \cdot \mathbf{u}_{(\alpha)}) \cdot \nabla)\mathbf{u} - (F\nabla \cdot \mathbf{u}_{(\alpha)})^\perp \text{curl } \mathbf{u}, J^t \mathbf{u}_{(\alpha)})_{L^2} \\ &= \frac{\mu}{2}(F \cdot \mathbf{u}_{(\alpha)}, \mathbf{u} \cdot (J^t \nabla \nabla \cdot \mathbf{u}_{(\alpha)}))_{L^2} - \frac{\mu}{2}(F(\mathbf{u} \cdot \nabla \nabla \cdot \mathbf{u}_{(\alpha)}), J^t \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + \frac{\mu}{2}(F \cdot \mathbf{u}_{(\alpha)}, (\nabla \cdot \mathbf{u})J^t \nabla \cdot \mathbf{u}_{(\alpha)})_{L^2} - \frac{\mu}{2}((\nabla \cdot \mathbf{u}_{(\alpha)})(\mathbf{u} \cdot \nabla)F, J^t \mathbf{u}_{(\alpha)})_{L^2} \\ &\quad - \frac{\mu}{2}(((F\nabla \cdot \mathbf{u}_{(\alpha)}) \cdot \nabla)\mathbf{u} - (F\nabla \cdot \mathbf{u}_{(\alpha)})^\perp \text{curl } \mathbf{u}, J^t \mathbf{u}_{(\alpha)})_{L^2}, \end{aligned}$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

where we used again the identity (4.13), and integration by parts. All the components are estimated as above by Cauchy-Schwarz inequality, continuous Sobolev embedding $H^2 \subset L^\infty$, (4.4) as well as (4.7) for the first line. It follows

$$\epsilon|B_3^t| \leq \mu C(h^*)F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{u}_{(\alpha)}|_{L^2}|\nabla \cdot \mathbf{u}_{(\alpha)}|_{L^2}. \quad (4.17)$$

One has

$$\begin{aligned} B_4^t &\stackrel{\text{def}}{=} \mu\beta^2((\mathbf{u} \cdot \nabla)(h(\nabla b \cdot \mathbf{u}_{(\alpha)})\nabla b), J^t\mathbf{u}_{(\alpha)})_{L^2} \\ &\quad + (h(\nabla b \cdot \mathbf{u}_{(\alpha)})(\nabla b \cdot \nabla)\mathbf{u} - h(\nabla b \cdot \mathbf{u}_{(\alpha)})(\nabla b)^\perp \text{curl } \mathbf{u}, J^t\mathbf{u}_{(\alpha)})_{L^2}. \end{aligned}$$

The second component is estimated by Cauchy-Schwarz inequality, continuous Sobolev embedding $H^2 \subset L^\infty$ and (4.4); and the first one after integrating by parts, averaging and repeated use of the commutator estimate (4.7). One obtains eventually

$$\epsilon|B_4^t| \leq \mu C(h^*)F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{u}_{(\alpha)}|_{L^2}^2. \quad (4.18)$$

We have one last term to estimate, namely

$$B_5^t \stackrel{\text{def}}{=} ((\mathbf{u} \cdot \nabla)\mathbf{v}_{(\alpha)} + (\mathbf{v}_{(\alpha)} \cdot \nabla)\mathbf{u} - \mathbf{v}_{(\alpha)}^\perp \text{curl } \mathbf{u}, [J^t, h]\mathbf{u}_{(\alpha)})_{L^2}.$$

By (4.7) and (2.4) in Lemma 2.2, and (2.2) in Lemma 2.1, one has

$$\begin{aligned} \epsilon|((\mathbf{v}_{(\alpha)} \cdot \nabla)\mathbf{u} - \mathbf{v}_{(\alpha)}^\perp \text{curl } \mathbf{u}, [J^t, h]\mathbf{u}_{(\alpha)})_{L^2}| \\ \leq F(\beta|\nabla b|_{H^2}, \epsilon|\nabla\zeta|_{H^2}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{v}_{(\alpha)}|_{Y^0}|\mathbf{u}_{(\alpha)}|_{L^2}. \end{aligned}$$

Then, one has, integrating by parts,

$$\begin{aligned} \epsilon|((\mathbf{u} \cdot \nabla)\mathbf{v}_{(\alpha)}, [J^t, h]\mathbf{u}_{(\alpha)})_{L^2}| \\ \leq \epsilon|(\mathbf{v}_{(\alpha)}, (\nabla \cdot \mathbf{u})[J^t, h]\mathbf{u}_{(\alpha)})_{L^2}| + \epsilon|(\mathbf{v}_{(\alpha)}, (\mathbf{u} \cdot \nabla)[J^t, h]\mathbf{u}_{(\alpha)})_{L^2}|. \end{aligned}$$

The first term is estimated as above, and the second by duality, since

$$\begin{aligned} |(\mathbf{u} \cdot \nabla)[J^t, h]\mathbf{u}_{(\alpha)}|_{X^0} &\lesssim \epsilon|(\mathbf{u} \cdot \nabla)[J^t, h]\mathbf{u}_{(\alpha)}|_{L^2} + \sqrt{\mu}\epsilon|\nabla \cdot ((\mathbf{u} \cdot \nabla)[J^t, h]\mathbf{u}_{(\alpha)})|_{L^2} \\ &\leq C(\mu)F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^3}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{u}_{(\alpha)}|_{X^0}, \end{aligned}$$

where we used (4.7) and continuous Sobolev embedding $H^2 \subset L^\infty$. Thus

$$\epsilon|B_5^t| \leq C(\mu)F(\beta|\nabla b|_{H^3}, \epsilon|\nabla\zeta|_{H^3}, \epsilon|\mathbf{u}|_{H^3})|\mathbf{v}_{(\alpha)}|_{Y^0}|\mathbf{u}_{(\alpha)}|_{L^2}. \quad (4.19)$$

Collecting estimates (4.14)–(4.19) and using Lemma 4.1, we deduce that

$$\epsilon B^t = \epsilon(B_0^t + B_1^t + B_2^t + B_3^t + B_4^t + B_5^t)$$

is estimated as in the statement. This concludes the proof. \square

5. Well-posedness

We are now in position to prove our main results concerning the Cauchy problem for (1.3).

Proposition 5.1 (Existence and uniqueness). *Let $N \geq 4$, $b \in \dot{H}^{N+2}$ and $(\zeta_0, \mathbf{v}_0) \in H^N \times Y^N$ satisfying (1.7) with $h_*, h^* > 0$, and $\operatorname{curl} \mathbf{v}_0 \in H^{N-1}$. Then there exists $T > 0$ and a unique $(\zeta, \mathbf{v}) \in L^\infty(0, T; H^N \times Y^N) \cap C([0, T]; H^2 \times (H^1)^d)$ satisfying (1.3) and $(\zeta, \mathbf{v})|_{t=0} = (\zeta_0, \mathbf{v}_0)$. Moreover, one can restrict*

$$T^{-1} = C(\mu, h_*^{-1}, h^*)F(\beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^4}, \epsilon|\mathbf{v}_0|_{Y^4}, \epsilon|\operatorname{curl} \mathbf{v}_0|_{H^3}) > 0$$

such that, for any $t \in [0, T]$, (1.7) holds with $\tilde{h}_* = h_*/2$, $\tilde{h}^* = 2h^*$, and

$$\mathcal{E}^N(\zeta, \mathbf{v}) + |\operatorname{curl} \mathbf{v}|_{H^{N-1}}^2 \leq \mathbf{C}_0 (\mathcal{E}^N(\zeta_0, \mathbf{v}_0) + |\operatorname{curl} \mathbf{v}_0|_{H^{N-1}}^2)$$

with $\mathbf{C}_0 = C(\mu, h_*^{-1}, h^*, \beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^4}, \epsilon|\mathbf{v}_0|_{Y^4}, \epsilon|\operatorname{curl} \mathbf{v}_0|_{H^3})$.

Proof. Construction. The construction of a solution is fairly classical, and follows the line of [32, Sec. 4.3.4]. Consider the mollified system (4.1) with right-hand side $r^\iota = 0$, $\mathbf{r}^\iota = \mathbf{0}$ and mollified initial data $(\zeta_0^\iota, \mathbf{v}_0^\iota) \stackrel{\text{def}}{=} (J^\iota \zeta_0, J^\iota \mathbf{v}_0)$. Obviously, for any $\iota \in (0, 1)$, $\zeta_0^\iota, \mathbf{v}_0^\iota \in H^n$ with arbitrary large $n \in \mathbb{N}$. By the Cauchy-Lipschitz theorem on Banach spaces, there exists a unique (smooth) solution to (4.1) with initial data $(\zeta_0^\iota, \mathbf{v}_0^\iota)$, that we denote $U^\iota \stackrel{\text{def}}{=} (\zeta^\iota, \mathbf{v}^\iota)$, defined on the maximal time interval $[0, T^\iota)$. For any multi-index $\alpha \in \mathbb{N}^d$, denote

$$\zeta_{(\alpha)}^\iota \stackrel{\text{def}}{=} \partial^\alpha \zeta^\iota \quad \text{and} \quad \mathbf{v}_{(\alpha)}^\iota \stackrel{\text{def}}{=} \partial^\alpha \mathbf{v}^\iota - \mu \epsilon \nabla (w^\iota \partial^\alpha \zeta^\iota)$$

where $w^\iota \stackrel{\text{def}}{=} -h^\iota \nabla \cdot \mathbf{u}^\iota + \beta \nabla b \cdot \mathbf{u}^\iota$, $h^\iota \stackrel{\text{def}}{=} 1 + \epsilon \zeta^\iota - \beta b$, $\mathbf{u}^\iota \stackrel{\text{def}}{=} \mathfrak{T}[h^\iota, \beta b]^{-1}(h^\iota \mathbf{v}^\iota)$. By proceeding exactly as in the proof of Proposition 3.1, one easily checks that U^ι satisfies the quasilinear mollified system (4.2) with

$$r_{(\alpha)}^\iota = J^\iota r_{(\alpha)}(\zeta^\iota, \mathbf{v}^\iota) \quad \text{and} \quad \mathbf{r}_{(\alpha)}^\iota = J^\iota \mathbf{r}_{(\alpha)}(\zeta^\iota, \mathbf{v}^\iota) + \mu \epsilon \nabla r_{(\alpha)}^{\prime \iota},$$

where $r_{(\alpha)}$ and $\mathbf{r}_{(\alpha)}$ are given in Proposition 3.1, and

$$r_{(\alpha)}^{\prime \iota} = -(\operatorname{Id} - J^\iota)(\zeta_{(\alpha)}^\iota \partial_t w^\iota) - [J^\iota, w^\iota] \partial^\alpha \nabla \cdot (h^\iota \mathbf{u}^\iota).$$

By using (2.2) in Lemma 2.1, one has

$$\mu \epsilon |\nabla r_{(\alpha)}^{\prime \iota}|_{Y^0} \lesssim \epsilon \sqrt{\mu} |r_{(\alpha)}^{\prime \iota}|_{L^2},$$

which, by (4.4) and (4.7) in Lemma 4.3 and using bounds obtained in the proof of Proposition 3.1, is easily seen to satisfy the same estimate as $\mathbf{r}_{(\alpha)}$ (this is a simplification with regards to the proof of [32, Sec. 4.3.4]; see Remark 4.26 therein). Thus we have, for any $1 \leq |\alpha| \leq N$,

$$|r'_{(\alpha)}|_{L^2} + |\mathbf{r}'_{(\alpha)}|_{Y^0} \leq \mathbf{F} \left(|\zeta^\iota|_{H^{|\alpha|}} + |\mathbf{v}^\iota|_{Y^{|\alpha|}} + |\operatorname{curl} \mathbf{v}^\iota|_{H^{|\alpha|-1}} \right)$$

with $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^{N+1}}, \epsilon|\nabla\zeta^\iota|_{H^3}, \epsilon|\mathbf{v}^\iota|_{Y^4}, \epsilon|\operatorname{curl} \mathbf{v}^\iota|_{H^3})$. It follows, applying Proposition 4.4 and Proposition 4.5,

$$\begin{aligned} \frac{d}{dt} \mathcal{F}^N(\zeta^\iota, \mathbf{v}^\iota) &\leq \mathbf{F} \mathcal{F}^N(\zeta^\iota, \mathbf{v}^\iota) \\ &\quad + \mathbf{F} \left(|\zeta^\iota|_{H^N} + |\mathbf{v}^\iota|_{Y^N} + |\operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}} \right) \mathcal{F}^N(\zeta^\iota, \mathbf{v}^\iota)^{1/2} \end{aligned}$$

with \mathbf{F} as above, using that by (4.1)₁, (4.4) in Lemma 4.3, Lemma 2.4 and (2.7),(2.8) in Lemma 2.2,

$$|\partial_t \zeta^\iota|_{H^3} \leq |h^\iota \mathbf{u}^\iota|_{H^4} \leq C(\mu, h_\star^{-1}, h^\star, \epsilon|\nabla\zeta^\iota|_{H^3}, \beta|\nabla b|_{H^3})|\mathbf{v}^\iota|_{Y^4}.$$

Notice that (1.7) propagates for large time since

$$1 + \epsilon\zeta^\iota - \beta b = 1 + \epsilon\zeta_0^\iota - \beta b + \epsilon \int_0^\iota \partial_t \zeta^\iota \geq h_\star - \epsilon \int_0^\iota |\partial_t \zeta|_{L^\infty}.$$

Finally, applying the operator curl to (4.1)₂ (recall $r^\iota = 0, \mathbf{r}^\iota = \mathbf{0}$), one has

$$\partial_t \operatorname{curl} \mathbf{v}^\iota + \epsilon J^\iota \nabla \cdot (\mathbf{u}^\iota \operatorname{curl} \mathbf{v}^\iota) = 0.$$

Proceeding as in Proposition 3.1 to show that $\operatorname{curl} \partial^\alpha \mathbf{v}^\iota$ satisfies the same conservation law up to a tame remainder term, testing against $\operatorname{curl} \partial^\alpha \mathbf{v}^\iota$ to deduce energy estimates and summing over $0 \leq |\alpha| \leq N - 1$ yields

$$\frac{d}{dt} |\operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}}^2 \lesssim \epsilon |\mathbf{u}^\iota|_{H^4} |\operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}}^2 + \epsilon |\operatorname{curl} \mathbf{v}^\iota|_{H^3} |\mathbf{u}^\iota|_{H^N} |\operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}}.$$

Altogether, using Gronwall-type estimates, Lemmata 4.2 and 4.4 and straightforward arguments (see for instance [32, p. 109] for details), one deduces that there exists $T > 0$ with

$$T^{-1} = C(\mu, h_\star^{-1}, h^\star)F(\beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^4}, \epsilon|\mathbf{v}_0|_{Y^4}, \epsilon|\operatorname{curl} \mathbf{v}_0|_{H^3}) > 0$$

such that for any $\iota \in (0, 1)$, $T^\iota > T$; and for any $t \in [0, T]$,

$$\mathcal{E}^N(\zeta^\iota, \mathbf{v}^\iota) + |\operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}}^2 \leq \mathbf{C}_0 (\mathcal{E}^N(\zeta_0, \mathbf{v}_0) + |\operatorname{curl} \mathbf{v}_0|_{H^{N-1}}^2) \quad (5.1)$$

with $\mathbf{C}_0 = C(\mu, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^4}, \epsilon|\mathbf{v}_0|_{Y^4}, \epsilon|\operatorname{curl} \mathbf{v}_0|_{H^3})$, and (1.7) holds for $h^\iota = 1 + \epsilon\zeta^\iota - \beta b$ with $\tilde{h}_\star = h_\star/2$ and $\tilde{h}^\star = 2h^\star$.

Notice that the time interval $[0, T]$ and energy estimates are uniform with respect to $\iota \in (0, 1)$. We shall prove below that the sequence $(\zeta^\iota, \mathbf{v}^\iota)$ defines a Cauchy sequence whose limit provides the desired solution.

Convergence. Denote $\underline{\zeta} \stackrel{\text{def}}{=} \zeta^{\iota^2} - \zeta^{\iota^1}$ and $\underline{\mathbf{v}} \stackrel{\text{def}}{=} \mathbf{v}^{\iota^1} - \mathbf{v}^{\iota^2}$. Then $(\underline{\zeta}, \underline{\mathbf{v}})$ satisfies

$$\begin{cases} \partial_t \underline{\zeta} + J^{\iota^2} \nabla \cdot (\underline{h} \underline{\mathbf{u}}) = J^{\iota^2} \underline{r} + (J^{\iota^1} - J^{\iota^2}) \bar{r}, \\ \partial_t \underline{\mathbf{v}} + J^{\iota^2} \left(\nabla \underline{\zeta} + \epsilon \underline{\mathbf{u}}^\perp \text{curl} \underline{\mathbf{v}} + \frac{\epsilon}{2} \nabla (|\underline{\mathbf{u}}|^2) - \mu \epsilon \nabla (\mathcal{R}[\underline{h}, \underline{\mathbf{u}}] + \mathcal{R}_b[\underline{h}, \beta b, \underline{\mathbf{u}}]) \right) \\ \qquad \qquad \qquad = J^{\iota^2} \underline{\mathbf{r}} + (J^{\iota^1} - J^{\iota^2}) \bar{\mathbf{r}}, \end{cases} \quad (5.2)$$

with notation $\underline{h} = 1 + \epsilon \underline{\zeta} - \beta b$ and $\underline{\mathbf{u}} \stackrel{\text{def}}{=} \mathfrak{T}[\underline{h}, \beta b]^{-1}(h \mathbf{v})$, and

$$\begin{aligned} \underline{r} &= \nabla \cdot (\underline{h} \underline{\mathbf{u}}) + \nabla \cdot (h^{\iota^1} \mathbf{u}^{\iota^1}) - \nabla \cdot (h^{\iota^2} \mathbf{u}^{\iota^2}), & \bar{r} &= \nabla \cdot (h^{\iota^1} \mathbf{u}^{\iota^1}), \\ \underline{\mathbf{r}} &= \epsilon (\underline{\mathbf{u}}^\perp \text{curl} \underline{\mathbf{v}} + (\mathbf{u}^{\iota^1})^\perp \text{curl} \mathbf{v}^{\iota^1} - (\mathbf{u}^{\iota^2})^\perp \text{curl} \mathbf{v}^{\iota^2}) + \frac{\epsilon}{2} \nabla (|\underline{\mathbf{u}}|^2 + |\mathbf{u}^{\iota^1}|^2 - |\mathbf{u}^{\iota^2}|^2) \\ &\quad - \mu \epsilon \nabla (\mathcal{R}[\underline{h}, \underline{\mathbf{u}}] + \mathcal{R}[h^{\iota^1}, \mathbf{u}^{\iota^1}] - \mathcal{R}[h^{\iota^2}, \mathbf{u}^{\iota^2}]) \\ &\quad - \mu \epsilon \nabla (\mathcal{R}_b[\underline{h}, \beta b, \underline{\mathbf{u}}] + \mathcal{R}_b[h^{\iota^1}, \beta b, \mathbf{u}^{\iota^1}] - \mathcal{R}_b[h^{\iota^2}, \beta b, \mathbf{u}^{\iota^2}]), \\ \bar{\mathbf{r}} &= \nabla \zeta^{\iota^1} + \epsilon (\mathbf{u}^{\iota^1})^\perp \text{curl} \mathbf{v}^{\iota^1} + \frac{\epsilon}{2} \nabla (|\mathbf{u}^{\iota^1}|^2) - \mu \epsilon \nabla (\mathcal{R}[h^{\iota^1}, \mathbf{u}^{\iota^1}] + \mathcal{R}_b[h^{\iota^1}, \beta b, \mathbf{u}^{\iota^1}]). \end{aligned}$$

By using the previously obtained energy estimates (5.1), Lemma 2.4 and Lemma 2.11, one has

$$|\underline{\mathbf{u}}|_{X^1} + |\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2}|_{X^1} \leq \mathbf{C}_0 \mathcal{E}^1(\underline{\zeta}, \underline{\mathbf{v}})^{1/2}$$

where we denote, here and thereafter,

$$\mathbf{C}_0 = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^{N+1}}, \epsilon |\zeta_0|_{H^4}, \epsilon |\mathbf{v}_0|_{Y^4}, \epsilon |\text{curl} \mathbf{v}_0|_{H^3}).$$

It follows, after straightforward computations and using Lemmata 2.1, 2.2 and (4.6) in Lemma 4.3,

$$|J^{\iota^2} \underline{r}|_{L^2} + |J^{\iota^2} \underline{\mathbf{r}}|_{Y^0} \leq \mathbf{C}_0 (\mathcal{E}^1(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} + |\text{curl} \underline{\mathbf{v}}|_{L^2}).$$

By (4.6) in Lemma 4.3, we have immediately

$$|(J^{\iota^1} - J^{\iota^2}) \bar{r}|_{L^2} + |(J^{\iota^1} - J^{\iota^2}) \bar{\mathbf{r}}|_{Y^0} \leq \mathbf{C}_0 |\iota^2 - \iota^1|.$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

In the same way, setting $1 \leq |\alpha| \leq 2$, then $\underline{\zeta}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta^{\iota^2} - \partial^\alpha \zeta^{\iota^1}$ and $\underline{\mathbf{v}}_{(\alpha)} \stackrel{\text{def}}{=} \mathbf{v}_{(\alpha)}^{\iota^2} - \mathbf{v}_{(\alpha)}^{\iota^1}$ satisfy

$$\begin{cases} \partial_t \underline{\zeta}_{(\alpha)} + J^{\iota^2} \left(\epsilon \nabla \cdot (\mathbf{u}^{\iota^2} \underline{\zeta}_{(\alpha)}) + \nabla \cdot (h^{\iota^2} \mathfrak{T}[h^{\iota^2}, \beta b]^{-1} (h^{\iota^2} \underline{\mathbf{v}}_{(\alpha)})) \right) \\ \qquad \qquad \qquad = J^{\iota^2} \underline{r}_{(\alpha)} + (J^{\iota^1} - J^{\iota^2}) \bar{r}_{(\alpha)}, \\ \partial_t \underline{\mathbf{v}}_{(\alpha)} + J^{\iota^2} \left(\nabla \underline{\zeta}_{(\alpha)} + \epsilon (\mathbf{u}^{\iota^2})^\perp \text{curl} \underline{\mathbf{v}}_{(\alpha)} + \epsilon \nabla (\mathbf{u}^{\iota^2} \cdot \underline{\mathbf{v}}_{(\alpha)}) \right) \\ \qquad \qquad \qquad = J^{\iota^2} \underline{\mathbf{r}}_{(\alpha)} + (J^{\iota^1} - J^{\iota^2}) \bar{\mathbf{r}}_{(\alpha)} + \mu \epsilon \nabla (r'_{(\alpha)}{}^{\iota^2} - r'_{(\alpha)}{}^{\iota^1}), \end{cases} \quad (5.3)$$

where

$$\begin{aligned} \underline{r}_{(\alpha)} &= \nabla \cdot (\epsilon (\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2}) \zeta_{(\alpha)}^{\iota^1}) + h^{\iota^1} \mathfrak{T}[h^{\iota^1}, \beta b]^{-1} (h^{\iota^1} \mathbf{v}_{(\alpha)}^{\iota^1}) - h^{\iota^2} \mathfrak{T}[h^{\iota^2}, \beta b]^{-1} (h^{\iota^2} \mathbf{v}_{(\alpha)}^{\iota^1}) \\ &\quad + r_{(\alpha)}(\zeta^{\iota^2}, \mathbf{v}^{\iota^2}) - r_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}), \\ \bar{r}_{(\alpha)} &= \epsilon \nabla \cdot (\mathbf{u}^{\iota^1} \zeta_{(\alpha)}^{\iota^1}) + \nabla \cdot (h^{\iota^1} \mathbf{u}_{(\alpha)}^{\iota^1}) - r_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}), \\ \underline{\mathbf{r}}_{(\alpha)} &= \epsilon (\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2})^\perp \text{curl} \mathbf{v}_{(\alpha)}^{\iota^1} + \epsilon \nabla ((\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2}) \cdot \mathbf{v}_{(\alpha)}^{\iota^1}) \\ &\quad + \mathbf{r}_{(\alpha)}(\zeta^{\iota^2}, \mathbf{v}^{\iota^2}) - \mathbf{r}_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}), \\ \bar{\mathbf{r}}_{(\alpha)} &= \epsilon (\mathbf{u}^{\iota^1})^\perp \text{curl} \mathbf{v}_{(\alpha)}^{\iota^1} + \epsilon \nabla (\mathbf{u}^{\iota^1} \cdot \mathbf{v}_{(\alpha)}^{\iota^1}) - \mathbf{r}_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}). \end{aligned}$$

In order to estimate the right-hand side, notice first that $|\alpha| + 2 \leq 4 \leq N$, one has by (5.1), (2.2) in Lemma 2.1, (2.4) in Lemma 2.2 and Lemma 2.4:

$$|\zeta_{(\alpha)}^{\iota^1}|_{H^2} + |\mathbf{v}_{(\alpha)}^{\iota^1}|_{Y^2} + |\mathbf{u}_{(\alpha)}^{\iota^1}|_{X^2} \leq \mathbf{C}_0.$$

Moreover, we already indicated $|\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2}|_{X^1} \leq \mathbf{C}_0 \mathcal{E}^1(\underline{\zeta}, \underline{\mathbf{v}})^{1/2}$ and by (3.4) in Proposition 3.1,

$$\begin{aligned} &|r_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}) - r_{(\alpha)}(\zeta^{\iota^2}, \mathbf{v}^{\iota^2})|_{L^2} + |\mathbf{r}_{(\alpha)}(\zeta^{\iota^1}, \mathbf{v}^{\iota^1}) - \mathbf{r}_{(\alpha)}(\zeta^{\iota^2}, \mathbf{v}^{\iota^2})|_{Y^0} \\ &\qquad \qquad \qquad \leq \mathbf{C}_0 (\mathcal{E}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} + |\text{curl} \underline{\mathbf{v}}|_{H^1}). \end{aligned}$$

It follows, by (4.4) in Lemma 4.3 and Lemmata 2.1, 2.2, 2.3 2.4 and 2.11,

$$|J^{\iota^2} \underline{r}_{(\alpha)}|_{L^2} + |J^{\iota^2} \underline{\mathbf{r}}_{(\alpha)}|_{Y^0} \leq \mathbf{C}_0 (\mathcal{E}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} + |\text{curl} \underline{\mathbf{v}}|_{H^1})$$

and, by (5.1) and (4.6) in Lemma 4.3,

$$|(J^{\iota^1} - J^{\iota^2}) \bar{r}_{(\alpha)}|_{L^2} + |(J^{\iota^1} - J^{\iota^2}) \bar{\mathbf{r}}_{(\alpha)}|_{Y^0} \leq \mathbf{C}_0 |\iota^2 - \iota^1|.$$

Following the same remark as above, we control the last contribution:

$$\mu \epsilon |\nabla r'_{(\alpha)}{}^{\iota^2} - \nabla r'_{(\alpha)}{}^{\iota^1}|_{Y^0} \leq \sqrt{\mu} \epsilon |r'_{(\alpha)}{}^{\iota^2} - r'_{(\alpha)}{}^{\iota^1}|_{L^2} \leq \mathbf{C}_0 (\mathcal{E}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} + |\iota^2 - \iota^1|).$$

Thus applying Proposition 4.4 to (5.2), Proposition 4.5 to (5.3) and adapting Lemma 4.2, one has

$$\frac{d}{dt} \mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) \leq \mathbf{C}_0 \left(\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2 + |\iota^2 - \iota^1| \mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} \right).$$

with the notation $\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) \stackrel{\text{def}}{=} \sum_{0 \leq |\alpha| \leq 2} \mathcal{F}[h^{\iota^2}, \beta b](\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)})$. Notice also the identity

$$\begin{aligned} \partial_t \operatorname{curl} \underline{\mathbf{v}} + \epsilon J^{\iota^2} \nabla \cdot (\mathbf{u}^{\iota^2} \operatorname{curl} \underline{\mathbf{v}}) &= J^{\iota^2} \nabla \cdot ((\mathbf{u}^{\iota^1} - \mathbf{u}^{\iota^2}) \operatorname{curl} \underline{\mathbf{v}}^{\iota^1}) \\ &\quad + \epsilon (J^{\iota^1} - J^{\iota^2}) \nabla \cdot (\mathbf{u}^{\iota^1} \operatorname{curl} \underline{\mathbf{v}}^{\iota^1}), \end{aligned}$$

so that, proceeding as above,

$$\frac{d}{dt} |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2 \leq \mathbf{C}_0 \left(\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2 + |\iota^2 - \iota^1| \mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} \right).$$

Applying Gronwall's Lemma and since, by (4.6) in Lemma 4.3, one has

$$|\zeta_0^{\iota^2} - \zeta_0^{\iota^1}|_{H^2} + |\mathbf{v}_0^{\iota^2} - \mathbf{v}_0^{\iota^1}|_{H^2} \lesssim |\iota^2 - \iota^1| \mathcal{E}^N(\zeta_0, \mathbf{v}_0)^{1/2},$$

we find that

$$\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1} \leq \mathbf{C}_0 |\iota^2 - \iota^1| (1+t) \exp(\mathbf{C}_0 t).$$

Slightly adapting the proof Lemma 4.2 and thanks to (5.1) and Lemma 2.4 and 2.11, we deduce

$$\begin{aligned} |\zeta^{\iota^2} - \zeta^{\iota^1}|_{H^2} + |\mathbf{v}^{\iota^2} - \mathbf{v}^{\iota^1}|_{Y^2} + |\operatorname{curl} \mathbf{v}^{\iota^2} - \operatorname{curl} \mathbf{v}^{\iota^1}|_{H^1} + |\mathbf{u}^{\iota^2} - \mathbf{u}^{\iota^1}|_{X^2} \\ \leq \mathbf{C}_0 |\iota^2 - \iota^1| (1+t) \exp(\mathbf{C}_0 t). \end{aligned}$$

The Cauchy sequences are strongly convergent in low regularity Banach spaces $C([0, T]; H^n)$, and are also bounded thus weakly convergent (up to a subsequence) in high regularity spaces by (5.1). By uniqueness of the limit, there exists $(\zeta, \mathbf{v}, w, \mathbf{u}, v) \in L^\infty(0, T; H^N \times Y^N \times H^{N-1} \times X^N \times H^N)$ such that (1.7) holds with $\tilde{h}_* = h_*/2$ and $\tilde{h}^* = 2h^*$, satisfying the desired energy estimates and, as $\iota \rightarrow 0$,

$$\sup_{t \in [0, T]} (|\zeta^\iota - \zeta|_{H^2} + |\mathbf{v}^\iota - \mathbf{v}|_{H^1} + |\operatorname{curl} \mathbf{v}^\iota - w|_{H^2} + |\mathbf{u}^\iota - \mathbf{u}|_{H^2} + |\nabla \cdot \mathbf{u}^\iota - v|_{H^2}) \rightarrow 0.$$

By uniqueness of the limit, $w = \operatorname{curl} \mathbf{v}$, $v = \nabla \cdot \mathbf{u}$ and $\mathbf{u} = \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v})$. The level of regularity in the above convergence result is sufficient to pass to the limit in (4.1), so that (ζ, \mathbf{v}) is a strong solution to (1.3). That it satisfies the desired initial data is guaranteed by (4.3) in Lemma 4.3.

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Uniqueness. By considering $\underline{\zeta}, \underline{\mathbf{v}}$ the difference between two solutions with same initial data, and proceeding exactly as above (with fewer terms since mollifications are not involved), we find

$$\frac{d}{dt} \left(\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2 \right) \leq \mathbf{C}_0 \left(\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2 \right).$$

Applying Gronwall's estimate and since $(\mathcal{F}^2(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^1}^2)|_{t=0} = 0$, we deduce $\underline{\zeta} = 0, \underline{\mathbf{v}} = \mathbf{0}$. This concludes the proof of Proposition 5.1. \square

Proposition 5.2 (Stability). *Let $N \geq 4$ and $(b, \zeta^0, \mathbf{v}^0)$ satisfy the assumptions of Proposition 5.1, and denote*

$$M_0 = |\zeta_0|_{H^N} + |\mathbf{v}_0|_{Y^N} + |\operatorname{curl} \mathbf{v}_0|_{H^N}$$

Let $1 \leq n \leq N$ and $(\tilde{\zeta}, \tilde{\mathbf{v}}, \operatorname{curl} \tilde{\mathbf{v}}) \in L^\infty(0, \tilde{T}; H^{N+1} \times Y^{N+1} \times H^N) \cap C([0, \tilde{T}]; H^n \times Y^n \times H^{n-1})$ satisfies (1.7) and (1.3) up to some remainders $(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}, \operatorname{curl} \tilde{\mathbf{r}}) \in L^1(0, \tilde{T}; H^n \times Y^n \times H^{n-1})$. Denote

$$\tilde{M} = \operatorname{ess\,sup}_{t \in [0, \tilde{T}]} (|\tilde{\zeta}|_{H^{N+1}} + |\tilde{\mathbf{v}}|_{Y^{N+1}} + |\operatorname{curl} \tilde{\mathbf{v}}|_{H^N}).$$

Then there exists

$$T^{-1} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^{N+1}}, \epsilon M_0, \epsilon \tilde{M}) > 0,$$

such that for any $t \in [0, \min(T, \tilde{T})]$,

$$\begin{aligned} & (|\zeta - \tilde{\zeta}|_{H^n} + |\mathbf{v} - \tilde{\mathbf{v}}|_{Y^n} + |\operatorname{curl} \mathbf{v} - \operatorname{curl} \tilde{\mathbf{v}}|_{H^{n-1}})(t) \leq \\ & \mathbf{C} (|\zeta - \tilde{\zeta}|_{H^n} + |\mathbf{v} - \tilde{\mathbf{v}}|_{Y^n} + |\operatorname{curl} \mathbf{v} - \operatorname{curl} \tilde{\mathbf{v}}|_{H^{n-1}})|_{t=0} \\ & + \mathbf{C} \int_0^t (|\tilde{\mathbf{r}}|_{H^n} + |\tilde{\mathbf{r}}|_{Y^n} + |\operatorname{curl} \tilde{\mathbf{r}}|_{H^{n-1}})(t') dt'. \end{aligned}$$

with $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^{N+1}}, \epsilon M_0, \epsilon \tilde{M})$.

Proof. This Lipschitz stability property was already at stake in the convergence part of the proof of Proposition 5.1. We use the same strategy.

Denote $\underline{\zeta} \stackrel{\text{def}}{=} \zeta - \tilde{\zeta}$, $\underline{\mathbf{v}} \stackrel{\text{def}}{=} \mathbf{v} - \tilde{\mathbf{v}}$, $\underline{h} \stackrel{\text{def}}{=} 1 + \epsilon \underline{\zeta} - \beta b$ and $\underline{\mathbf{u}} \stackrel{\text{def}}{=} \mathfrak{T}[\underline{h}, \beta b]^{-1}(\underline{h} \underline{\mathbf{v}})$; as well as $\underline{\zeta}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta - \partial^\alpha \tilde{\zeta}$, $\underline{\mathbf{v}}_{(\alpha)} \stackrel{\text{def}}{=} \mathbf{v}_{(\alpha)} - \tilde{\mathbf{v}}_{(\alpha)}$ and $\underline{\mathbf{u}}_{(\alpha)} \stackrel{\text{def}}{=} \mathfrak{T}[\underline{h}, \beta b]^{-1}(\underline{h} \underline{\mathbf{v}}_{(\alpha)})$. Consider the system satisfied by $(\underline{\zeta}, \underline{\mathbf{v}})$ and $(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)})$. Compared with (5.2)

and (5.3), there are fewer terms in the right-hand side, but they require more precise estimates. For $1 \leq |\alpha| \leq n$, one has

$$\left\{ \begin{array}{l} \partial_t \underline{\zeta}_{(\alpha)} + \epsilon \nabla \cdot (\mathbf{u} \underline{\zeta}_{(\alpha)}) + \nabla \cdot (h \mathbf{u}_{(\alpha)}) \\ \qquad \qquad \qquad = \underline{r}_{(\alpha)} - \partial^\alpha \tilde{r} + r_{(\alpha)}(\zeta, \mathbf{v}) - r_{(\alpha)}(\tilde{\zeta}, \tilde{\mathbf{v}}) \\ \\ \partial_t \underline{\mathbf{v}}_{(\alpha)} + \nabla \underline{\zeta}_{(\alpha)} + \epsilon \mathbf{u}^\perp \operatorname{curl} \underline{\mathbf{v}}_{(\alpha)} + \epsilon \nabla (\mathbf{u} \cdot \underline{\mathbf{v}}_{(\alpha)}) \\ \qquad \qquad \qquad = \underline{\mathbf{r}}_{(\alpha)} - \partial^\alpha \tilde{\mathbf{r}} + \mu \epsilon \nabla (\tilde{w} \partial^\alpha \tilde{r}) + \mathbf{r}_{(\alpha)}(\zeta, \mathbf{v}) - \mathbf{r}_{(\alpha)}(\tilde{\zeta}, \tilde{\mathbf{v}}) \end{array} \right. \quad (5.4)$$

where $\tilde{w} \stackrel{\text{def}}{=} -\tilde{h} \nabla \cdot \tilde{\mathbf{u}} + \beta \nabla b \cdot \tilde{\mathbf{u}}$, $\tilde{h} \stackrel{\text{def}}{=} 1 + \epsilon \tilde{\zeta} - \beta b$, $\tilde{\mathbf{u}} \stackrel{\text{def}}{=} \mathfrak{T}[\tilde{h}, \beta b]^{-1}(\tilde{h} \tilde{\mathbf{v}})$, and

$$\begin{aligned} \underline{r}_{(\alpha)} &= \nabla \cdot (\epsilon (\tilde{\mathbf{u}} - \mathbf{u}) \tilde{\zeta}_{(\alpha)} + \tilde{h} \mathfrak{T}[\tilde{h}, \beta b]^{-1}(\tilde{h} \tilde{\mathbf{v}}_{(\alpha)}) - h \mathfrak{T}[h, \beta b]^{-1}(h \tilde{\mathbf{v}}_{(\alpha)})), \\ \underline{\mathbf{r}}_{(\alpha)} &= \epsilon (\tilde{\mathbf{u}} - \mathbf{u})^\perp \operatorname{curl} \tilde{\mathbf{v}}_{(\alpha)} + \epsilon \nabla ((\tilde{\mathbf{u}} - \mathbf{u}) \cdot \tilde{\mathbf{v}}_{(\alpha)}). \end{aligned}$$

Since $|\alpha| \leq N$, one has by (2.2) in Lemma 2.1 and Lemmata 2.2 and 2.4 to estimate $|\tilde{w}|_{H^2}$,

$$\begin{aligned} \epsilon |\tilde{\zeta}_{(\alpha)}|_{H^1} + \epsilon |\tilde{\mathbf{v}}_{(\alpha)}|_{Y^1} + \epsilon |\operatorname{curl} \tilde{\mathbf{v}}_{(\alpha)}|_{L^2} \\ \leq C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^3}, \epsilon |\tilde{\zeta}|_{H^{N+1}}, \epsilon |\tilde{\mathbf{v}}|_{Y^{N+1}}, \epsilon |\operatorname{curl} \tilde{\mathbf{v}}|_{H^N}). \end{aligned}$$

Using Lemmata 2.1, 2.2, 2.4, and 2.11, one checks that for $m \in \{0, 1, 2, 3\}$,

$$|\mathbf{u} - \tilde{\mathbf{u}}|_{H^m} \leq |\mathbf{u} - \tilde{\mathbf{u}}|_{X^m} \leq \mathbf{C} \times (|\underline{\zeta}|_{H^m} + |\underline{\mathbf{v}}|_{Y^m})$$

with $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^3}, \epsilon |\zeta|_{H^3}, \epsilon |\mathbf{v}|_{Y^3}, \epsilon |\tilde{\zeta}|_{H^3}, \epsilon |\tilde{\mathbf{v}}|_{Y^4})$. It follows, using again Lemmata 2.1, 2.2, 2.4, and adapting the proof of Lemma 2.11 to replace $|\mathbf{v}|_{Y^{2\nu n}} |\zeta - \tilde{\zeta}|_{H^n}$ with $|\mathbf{v}|_{Y^n} |\zeta - \tilde{\zeta}|_{H^{2\nu n}}$,

$$|\underline{r}_{(\alpha)}|_{L^2} + |\underline{\mathbf{r}}_{(\alpha)}|_{Y^0} \leq \epsilon \mathbf{C} (|\underline{\zeta}|_{H^3} + |\underline{\mathbf{v}}|_{Y^3}) (|\tilde{\zeta}_{(\alpha)}|_{H^1} + |\tilde{\mathbf{v}}_{(\alpha)}|_{Y^1} + |\operatorname{curl} \tilde{\mathbf{v}}_{(\alpha)}|_{L^2}),$$

with \mathbf{C} as above. Moreover, by (3.4) in Proposition 3.1, one has

$$\begin{aligned} |r_{(\alpha)}(\zeta, \mathbf{v}) - r_{(\alpha)}(\tilde{\zeta}, \tilde{\mathbf{v}})|_{L^2} + |\mathbf{r}_{(\alpha)}(\zeta, \mathbf{v}) - \mathbf{r}_{(\alpha)}(\tilde{\zeta}, \tilde{\mathbf{v}})|_{Y^0} \\ \leq \tilde{\mathbf{F}} \left(|\underline{\zeta}|_{H^{|\alpha|}} + |\underline{\mathbf{v}}|_{Y^{|\alpha|}} + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{|\alpha|-1}} \right) \end{aligned}$$

with

$$\begin{aligned} \tilde{\mathbf{F}} &= C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^{4\nu|\alpha|+1}}, \epsilon |\zeta|_{H^{4\nu|\alpha|}}, \epsilon |\mathbf{v}|_{Y^{4\nu|\alpha|}}, \epsilon |\operatorname{curl} \mathbf{v}|_{H^{3\nu|\alpha|-1}}, \\ &\quad \epsilon |\tilde{\zeta}|_{H^{4\nu|\alpha|}}, \epsilon |\tilde{\mathbf{v}}|_{Y^{4\nu|\alpha|}}, \epsilon |\operatorname{curl} \tilde{\mathbf{v}}|_{H^{3\nu|\alpha|-1}}). \end{aligned}$$

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

Finally, using (2.3) in Lemma 2.1, one finds as above

$$|\mu \epsilon \nabla(\tilde{w} \partial^\alpha \tilde{r})|_{Y^0} \leq C(\mu) |\epsilon \tilde{w} \partial^\alpha \tilde{r}|_{L^2} \leq \mathbf{C} |\partial^\alpha \tilde{r}|_{L^2}.$$

By Proposition 4.5, the energy estimate of Proposition 5.1 and using the identity $\partial_t \zeta = -\nabla \cdot (h \mathbf{u})$ to control $|\partial_t \zeta|_{H^3}$, we obtain

$$\begin{aligned} \frac{d}{dt} \mathcal{F}(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)}) &\leq \mathbf{F}(\mathcal{F}(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)}) + \mathcal{E}^{3\nu|\alpha|}(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{|\alpha|-1}}^2) \\ &\quad + \mathbf{C}(\mathcal{F}(\partial^\alpha \tilde{r}, \partial^\alpha \tilde{\mathbf{r}}) \mathcal{F}(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)}))^{1/2}, \end{aligned}$$

where we denote $\mathbf{F} = C(\mu, h_\star^{-1}, h^\star) F(\beta |\nabla b|_{H^{N+1}}, \epsilon M_0, \epsilon \tilde{M})$, and $\mathbf{C} = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^{N+1}}, \epsilon M_0, \epsilon \tilde{M})$.

In order to control $\operatorname{curl} \underline{\mathbf{v}}$, we notice that

$$\partial_t \operatorname{curl} \underline{\mathbf{v}} + \epsilon \nabla \cdot (\mathbf{u} \operatorname{curl} \underline{\mathbf{v}}) = \epsilon \nabla \cdot ((\tilde{\mathbf{u}} - \mathbf{u}) \operatorname{curl} \tilde{\mathbf{v}}) - \operatorname{curl} \tilde{\mathbf{r}},$$

so that standard energy estimates yield

$$\begin{aligned} \frac{d}{dt} (|\operatorname{curl} \underline{\mathbf{v}}|_{H^{n-1}}^2) &\lesssim |\operatorname{curl} \underline{\mathbf{v}}|_{H^{n-1}} \\ &\quad \times \left(\epsilon |\mathbf{u}|_{H^{2\nu n}} |\operatorname{curl} \underline{\mathbf{v}}|_{H^{n-1}} + \epsilon |\tilde{\mathbf{u}} - \mathbf{u}|_{H^n} |\operatorname{curl} \tilde{\mathbf{v}}|_{H^{2\nu n}} + |\operatorname{curl} \tilde{\mathbf{r}}|_{H^{n-1}} \right). \end{aligned}$$

Finally, one easily checks that, as in Lemma 4.2, that

$$\mathcal{E}^n(\underline{\zeta}, \underline{\mathbf{v}}) \leq \mathbf{C} \sum_{0 \leq |\alpha| \leq n} \mathcal{F}(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)}); \quad \sum_{0 \leq |\alpha| \leq n} \mathcal{F}(\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)}) \leq \mathbf{C} \mathcal{E}^n(\underline{\zeta}, \underline{\mathbf{v}}).$$

Thus for any $n \geq 3$, adding the above energy estimates for $1 \leq |\alpha| \leq n$, the corresponding one based on Proposition 4.4 when $\alpha = (0, 0)$, and by Gronwall's Lemma, we find

$$\begin{aligned} |\underline{\zeta}|_{H^n} + |\underline{\mathbf{v}}|_{Y^n} + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{n-1}} &\leq (|\underline{\zeta}|_{t=0}|_{H^n} + |\underline{\mathbf{v}}|_{t=0}|_{Y^n} + |\operatorname{curl} \underline{\mathbf{v}}|_{t=0}|_{H^{n-1}}) e^{\mathbf{F}t} \\ &\quad + \mathbf{C} \int_0^t (|\tilde{\mathbf{r}}|_{H^n} + |\tilde{\mathbf{r}}|_{Y^n} + |\operatorname{curl} \tilde{\mathbf{r}}|_{H^{n-1}})(t') e^{\mathbf{F}(t-t')} dt'. \end{aligned}$$

with \mathbf{C}, \mathbf{F} as above. The proposition is proved for $n \geq 3$. The case $n \leq 2$ is obtained in the same way, but using the estimates

$$\begin{aligned} |\underline{\mathbf{r}}_{(\alpha)}|_{L^2} + |\underline{\mathbf{r}}_{(\alpha)}|_{Y^0} &\leq \epsilon \mathbf{C} (|\underline{\zeta}|_{H^1} + |\underline{\mathbf{v}}|_{Y^1}) (|\tilde{\zeta}_{(\alpha)}|_{H^3} + |\tilde{\mathbf{v}}_{(\alpha)}|_{H^3} + |\operatorname{curl} \tilde{\mathbf{v}}_{(\alpha)}|_{H^2}) \\ &\quad \text{(notice that in that case, } |\alpha| \leq 2 \leq N + 1 - 3). \quad \square \end{aligned}$$

We now conclude this section with continuity results, completing the well-posedness of the Cauchy problem for system (1.3) in the sense of Hadamard.

Proposition 5.3 (Well-posedness). *Under the hypotheses of Proposition 5.1, the unique strong solution to (1.3) satisfies additionally $(\zeta, \mathbf{v}, \operatorname{curl} \mathbf{v}) \in C([0, T]; H^N \times Y^N \times H^{N-1})$. Moreover, the mapping $(\zeta_0, \mathbf{v}_0, \operatorname{curl} \mathbf{v}_0) \in H^N \times Y^N \times H^{N-1} \mapsto (\zeta, \mathbf{v}, \operatorname{curl} \mathbf{v}) \in C([0, T]; H^N \times Y^N \times H^{N-1})$ is continuous.*

More precisely, for $(\zeta_0, \mathbf{v}_0, \operatorname{curl} \mathbf{v}_0) \in H^N \times Y^N \times H^{N-1}$ satisfying (1.7) and $(\zeta_{0,n}, \mathbf{v}_{0,n}, \operatorname{curl} \mathbf{v}_{0,n}) \rightarrow (\zeta_0, \mathbf{v}_0, \operatorname{curl} \mathbf{v}_0)$ in $H^N \times Y^N \times H^{N-1}$, one can set $T^{-1} = C(\mu, h_\star^{-1}, h^\star) \times F(\beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^N}, \epsilon|\mathbf{v}_0|_{Y^N}, |\operatorname{curl} \mathbf{v}_0|_{H^{N-1}})$ and $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, there exists a unique strong solution $(\zeta_n, \mathbf{v}_n) \in C([0, T]; H^N \times Y^N)$ satisfying (1.3) with initial data $(\zeta_n, \mathbf{v}_n)|_{t=0} = (\zeta_{0,n}, \mathbf{v}_{0,n})$, and one has

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \left(|\zeta_n - \zeta|_{H^N} + |\mathbf{v}_n - \mathbf{v}|_{Y^N} + |\operatorname{curl} \mathbf{v}_n - \operatorname{curl} \mathbf{v}|_{H^{N-1}} \right) = 0.$$

Proof. Our proof is based on the Bona-Smith technique [9]. For $\iota \in (0, 1)$, denote $(\zeta^\iota, \mathbf{v}^\iota)$ (resp. $(\zeta_n^\iota, \mathbf{v}_n^\iota)$) the unique solution to (1.3) with mollified initial data $(\zeta_0^\iota, \mathbf{v}_0^\iota) \stackrel{\text{def}}{=} (J^\iota \zeta_0, J^\iota \mathbf{v}_0)$ (resp. $(\zeta_{0,n}^\iota, \mathbf{v}_{0,n}^\iota) \stackrel{\text{def}}{=} (J^\iota \zeta_{0,n}, J^\iota \mathbf{v}_{0,n})$) in $C([0, T]; H^2 \times (H^1)^d) \cap L^\infty(0, T; H^N \times Y^N)$, as provided by Proposition 5.1; see Lemma 4.3 for the definition of J^ι and relevant properties. In particular, by (4.4), one can restrict $n_0 \in \mathbb{N}$ such that the energy estimate and lower bound on T stated in Proposition 5.1 hold uniformly with respect to $\iota \in (0, 1)$ and $n \geq n_0$.

We then proceed as in the proof of Proposition 5.2, with $(\tilde{\zeta}, \tilde{\mathbf{v}}) = (\zeta^\iota, \mathbf{v}^\iota)$ and $(r, \mathbf{r}) = \mathbf{0}$. We thus find that the difference, $\underline{\zeta} \stackrel{\text{def}}{=} \zeta - \zeta^\iota$ and $\underline{\mathbf{v}} = \mathbf{v} - \mathbf{v}^\iota$ satisfies, for any $3 \leq m \leq N$,

$$\begin{aligned} \frac{d}{dt} (\mathcal{F}^m(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{m-1}}^2) &\leq \mathbf{C}_0 (\mathcal{F}^m(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{m-1}}^2) \\ &\quad + \epsilon \mathbf{C}_0 \mathcal{F}^m(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} \mathcal{E}^3(\underline{\zeta}, \underline{\mathbf{v}})^{1/2} (|\zeta^\iota|_{H^{m+1}} + |\mathbf{v}^\iota|_{Y^{m+1}} + |\operatorname{curl} \mathbf{v}^\iota|_{H^m}), \end{aligned}$$

with $\mathcal{F}^m(\underline{\zeta}, \underline{\mathbf{v}}) \stackrel{\text{def}}{=} \sum_{0 \leq |\alpha| \leq m} \mathcal{F}[h, \beta b](\underline{\zeta}_{(\alpha)}, \underline{\mathbf{v}}_{(\alpha)})$ where $\underline{\zeta}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta - \partial^\alpha \tilde{\zeta}$, $\underline{\mathbf{v}}_{(\alpha)} \stackrel{\text{def}}{=} \mathbf{v}_{(\alpha)} - \tilde{\mathbf{v}}_{(\alpha)}$, and (using (4.4) in Lemma 4.3 and the energy estimate in Proposition 5.1)

$$\mathbf{C}_0 = C(\mu, h_\star^{-1}, h^\star, \beta|\nabla b|_{H^{N+1}}, \epsilon|\zeta_0|_{H^N}, \epsilon|\mathbf{v}_0|_{Y^N}, \epsilon|\operatorname{curl} \mathbf{v}_0|_{H^{N-1}}).$$

By (4.5) in Lemma 4.3 and the energy estimate in Proposition 5.1, we find

$$|\zeta^\iota|_{H^{N+1}} + |\mathbf{v}^\iota|_{Y^{N+1}} + |\operatorname{curl} \mathbf{v}^\iota|_{H^N} \leq \iota^{-1} \mathbf{C}_0.$$

Moreover, since $3 + 1 \leq N$, the above differential energy inequality with $m = 3$ reads

$$\frac{d}{dt} (\mathcal{F}^N(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{N-1}}^2) \leq \mathbf{C}_0 (\mathcal{F}^N(\underline{\zeta}, \underline{\mathbf{v}}) + |\operatorname{curl} \underline{\mathbf{v}}|_{H^{N-1}}^2 + \mathcal{E}^3(\underline{\zeta}, \underline{\mathbf{v}})),$$

so that, adapting Lemma 4.2 and Gronwall's Lemma, and finally applying (4.3) in Lemma 4.3,

$$\sup_{t \in [0, T]} \iota^{-1} \mathcal{E}^3(\underline{\zeta}, \underline{\mathbf{v}}) \leq \mathbf{C}_0 \exp(\mathbf{C}_0 T) \times \iota^{-1} \mathcal{E}^3(\underline{\zeta}_0, \underline{\mathbf{v}}_0) \rightarrow 0 \quad (\iota \rightarrow 0).$$

Thus applying Gronwall's Lemma to the differential energy inequality with $m = N$ and again adapting Lemma 4.2 yields

$$\lim_{\iota \rightarrow 0} \sup_{t \in [0, T]} (|\zeta - \zeta^\iota|_{H^N} + |\mathbf{v} - \mathbf{v}^\iota|_{Y^N} + |\operatorname{curl} \mathbf{v} - \operatorname{curl} \mathbf{v}^\iota|_{H^{N-1}}) = 0,$$

Using that, for any $\iota \in (0, 1)$, $(\zeta^\iota, \mathbf{v}^\iota) \in C([0, T]; H^N \times Y^N)$ (by the smoothness of the initial data, Proposition 5.1, and integrating (1.3) with respect to time), we deduce $(\zeta, \mathbf{v}) \in C([0, T]; H^N \times Y^N)$.

Now we turn to the continuity of the flow map. The proof above yields

$$\lim_{\iota \rightarrow 0} \sup_{t \in [0, T]} (|\zeta_n - \zeta_n^\iota|_{H^N} + |\mathbf{v}_n - \mathbf{v}_n^\iota|_{Y^N} + |\operatorname{curl} \mathbf{v}_n - \operatorname{curl} \mathbf{v}_n^\iota|_{H^{N-1}}) = 0$$

uniformly with respect to $n \geq n_0$ (notice in particular the uniformity with respect to n in Lemma 4.3). Moreover, proceeding in the same way, we find the following estimate for $(\underline{\zeta}^\iota, \underline{\mathbf{v}}^\iota) \stackrel{\text{def}}{=} (\zeta_n^\iota - \zeta^\iota, \mathbf{v}_n^\iota - \mathbf{v}^\iota)$ with any given $\iota \in (0, 1)$:

$$\begin{aligned} & \frac{d}{dt} (\mathcal{F}^N(\underline{\zeta}^\iota, \underline{\mathbf{v}}^\iota) + |\operatorname{curl} \underline{\mathbf{v}}^\iota|_{H^{N-1}}^2) \\ & \leq \mathbf{C}_0 (\mathcal{F}^N(\underline{\zeta}^\iota, \underline{\mathbf{v}}^\iota) + |\operatorname{curl} \underline{\mathbf{v}}^\iota|_{H^{N-1}}^2 + \iota^{-1} \mathcal{F}^N(\underline{\zeta}^\iota, \underline{\mathbf{v}}^\iota)^{1/2} \mathcal{E}^3(\underline{\zeta}^\iota, \underline{\mathbf{v}}^\iota)^{1/2}). \end{aligned}$$

By Gronwall's Lemma, adapting Lemma 4.2 and (4.4) in Lemma 4.3, we immediately deduce

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} (|\zeta^\iota - \zeta_n^\iota|_{H^N} + |\mathbf{v}^\iota - \mathbf{v}_n^\iota|_{Y^N} + |\operatorname{curl} \mathbf{v}^\iota - \operatorname{curl} \mathbf{v}_n^\iota|_{H^{N-1}}) = 0.$$

The continuity of the flow map follows from the above limits and triangular inequality. \square

6. Proof of the main results

We proved in the previous section the well-posedness and stability of the Cauchy problem for system (1.3). We show in this section how to transcribe these results to the original formulation of the Green-Naghdi system (1.1). It is claimed in [12, 39] that (1.3) and (1.1) are equivalent, after “judiciously differentiating by parts” and “lengthy calculations”. Since this fact is of considerable importance in our work and is rather tedious to check, we detail the calculations below. We then conclude this section with the proof of Theorems 1.1 and 1.2.

Proposition 6.1. *Let $b \in \dot{H}^5(\mathbb{R}^d)$ and $\zeta \in C^0([0, T]; H^4(\mathbb{R}^d))$ be such that (1.7) holds.*

Assume $\mathbf{u} \in C^0([0, T]; X^4)$ is such that (ζ, \mathbf{u}) satisfies (1.1). Then $\mathbf{v} \stackrel{\text{def}}{=} h^{-1}\mathfrak{T}[h, \beta b]\mathbf{u} \in C^0([0, T]; Y^4)$ is uniquely defined and (ζ, \mathbf{v}) satisfies (1.3).

Assume $\mathbf{v} \in C^0([0, T]; Y^4)$ is such that (ζ, \mathbf{v}) satisfies (1.3), then $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v}) \in C^0([0, T]; X^4)$ is uniquely defined and (ζ, \mathbf{u}) satisfies (1.1).

Proof. By Lemmata 2.1, 2.2 and 2.5, one easily checks that for any given $\mathbf{u} \in C^0([0, T]; X^4)$ and $\zeta \in C^0([0, T]; H^4(\mathbb{R}^d))$ such that (1.7) holds, then $\mathbf{v} \stackrel{\text{def}}{=} h^{-1}\mathfrak{T}[h, \beta b]\mathbf{u} \in C^0([0, T]; Y^4)$. Conversely, given $\mathbf{v} \stackrel{\text{def}}{=} \in C^0([0, T]; Y^4)$, that $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v})$ is well-defined and satisfies $\mathbf{u} \in C^0([0, T]; X^4)$ follows by Lemmata 2.3 and 2.4. The regularity of time derivatives is provided by the equations (1.1) or (1.3). This regularity is sufficient to ensure that all the identities below hold in, say, $L^2(\mathbb{R}^d)$.

Let us first notice that, by the identity $\frac{1}{2}\nabla(|\mathbf{u}|^2) = (\mathbf{u} \cdot \nabla)\mathbf{u} - \mathbf{u}^\perp \text{curl } \mathbf{u}$, all terms of order $\mathcal{O}(\mu)$ in (1.3) and (1.1) agree. Notice also that, as pointed out in Appendix B, system (1.3) can be rewritten as (B.11). It follows that to complete the proof, we only need to show that

$$\begin{aligned} [\partial_t, \mathcal{T}[h, \beta b]]\mathbf{u} + \epsilon\mathbf{u}^\perp \text{curl}(\mathcal{T}[h, \beta b]\mathbf{u}) + \epsilon\nabla(\mathbf{u} \cdot \mathcal{T}[h, \beta b]\mathbf{u} - \frac{1}{2}w^2) \\ = \epsilon\mathcal{Q}[h, \mathbf{u}] + \epsilon\mathcal{Q}_b[h, \beta b, \mathbf{u}], \end{aligned} \quad (6.1)$$

with $w \stackrel{\text{def}}{=}} (\beta\nabla b) \cdot \mathbf{u} - h\nabla \cdot \mathbf{u}$. We clarify below why (6.1) holds.

Let us first decompose

$$\begin{aligned}\mathcal{T}[h, \beta b]\mathbf{u} &= \frac{1}{h}\nabla\left(\frac{-h^3}{3}\nabla\cdot\mathbf{u} + \frac{\beta}{2}h^2\nabla b\cdot\mathbf{u}\right) + \beta\left(-\frac{1}{2}h\nabla\cdot\mathbf{u} + \beta(\nabla b\cdot\mathbf{u})\right)\nabla b \\ &\stackrel{\text{def}}{=} \frac{1}{h}\nabla f_1 + \beta f_2\nabla b.\end{aligned}$$

We use the identity valid for sufficiently regular scalar functions f, g :

$$\begin{aligned}\mathbf{u}^\perp\text{curl}(f\nabla g) + \nabla(\mathbf{u}\cdot(f\nabla g)) &= (\mathbf{u}\cdot\nabla)(f\nabla g) + f(\nabla g\cdot\nabla)\mathbf{u} - (\nabla g)^\perp\text{curl}(\mathbf{u}) \\ &= (\mathbf{u}\cdot\nabla f)\nabla g + f\nabla(\nabla g\cdot\mathbf{u})\end{aligned}$$

to deduce

$$\begin{aligned}\mathbf{u}^\perp\text{curl}(\mathcal{T}[h, \beta b]\mathbf{u}) + \nabla(\mathbf{u}\cdot\mathcal{T}[h, \beta b]\mathbf{u}) \\ = \frac{-1}{h^2}(\mathbf{u}\cdot\nabla h)\nabla f_1 + \frac{1}{h}\nabla(\nabla f_1\cdot\mathbf{u}) + \beta(\mathbf{u}\cdot\nabla f_2)\nabla b + \beta f_2\nabla(\nabla b\cdot\mathbf{u}).\end{aligned}\tag{6.2}$$

Now, using that b is time-independent and replacing $\partial_t\zeta = -\nabla\cdot(h\mathbf{u})$,

$$\begin{aligned}[\partial_t, \mathcal{T}[h, \beta b]]\mathbf{u} &= \frac{\epsilon\nabla\cdot(h\mathbf{u})}{h^2}\nabla f_1 + \frac{\epsilon}{h}\nabla((h\nabla\cdot(h\mathbf{u}))(h\nabla\cdot\mathbf{u} - \beta\mathbf{u}\cdot\nabla b)) \\ &\quad + \frac{\epsilon\beta}{2}(\nabla\cdot(h\mathbf{u}))(\nabla\cdot\mathbf{u})\nabla b.\end{aligned}\tag{6.3}$$

By (6.2) and (6.3), the desired identity (6.1) becomes

$$\begin{aligned}\frac{\nabla\cdot\mathbf{u}}{h}\nabla f_1 + \frac{1}{h}\nabla\left((\nabla f_1 + (h^2\nabla\cdot\mathbf{u})\nabla h - \beta h\nabla\cdot(h\mathbf{u})\nabla b)\cdot\mathbf{u} + h^3(\nabla\cdot\mathbf{u})^2\right) \\ + \frac{\beta}{2}(\nabla\cdot(h\mathbf{u}))(\nabla\cdot\mathbf{u})\nabla b + \beta(\mathbf{u}\cdot\nabla f_2)\nabla b + \beta f_2\nabla(\nabla b\cdot\mathbf{u}) - \frac{1}{2}\nabla((\beta\mathbf{u}\cdot\nabla b - h\nabla\cdot\mathbf{u})^2) \\ = \mathcal{Q}[h, \mathbf{u}] + \mathcal{Q}_b[h, \beta b, \mathbf{u}].\end{aligned}\tag{6.4}$$

The non-topographical contributions in (6.4) (*i.e.* setting $\beta = 0$, including in f_1) are easily seen to match:

$$\begin{aligned}\frac{-1}{3}\frac{\nabla\cdot\mathbf{u}}{h}\nabla(h^3\nabla\cdot\mathbf{u}) + \frac{1}{h}\nabla\left(\frac{-1}{3}h^3(\nabla(\nabla\cdot\mathbf{u}))\cdot\mathbf{u} + h^3(\nabla\cdot\mathbf{u})^2\right) - \frac{1}{2}\nabla((h\nabla\cdot\mathbf{u})^2) \\ = \frac{-1}{3h}\nabla\left(h^3((\mathbf{u}\cdot\nabla)(\nabla\cdot\mathbf{u}) - (\nabla\cdot\mathbf{u})^2)\right) = \mathcal{Q}[h, \mathbf{u}].\end{aligned}$$

The remaining contributions in (6.4) are more involved. Denoting $f = \nabla \cdot \mathbf{u}$ and $g = \beta \nabla b \cdot \mathbf{u}$,

$$\begin{aligned}
 & \frac{1}{2} \frac{f}{h} \nabla (h^2 g) + \frac{1}{h} \nabla \left(\frac{1}{2} \mathbf{u} \cdot \nabla (h^2 g) - h(\mathbf{u} \cdot \nabla h + hf)g \right) \\
 & \quad + \frac{\beta}{2} (\mathbf{u} \cdot \nabla h + hf) f \nabla b + \beta (\mathbf{u} \cdot \nabla (g - \frac{1}{2} hf)) \nabla b \\
 & \quad + (g - \frac{1}{2} hf) \nabla g - \frac{1}{2} \nabla (g^2 - 2ghf) \\
 & = \frac{1}{2} \frac{f}{h} \nabla (h^2 g) + \frac{1}{h} \nabla \left(\frac{1}{2} h^2 \mathbf{u} \cdot \nabla g - h^2 f g \right) + \frac{\beta}{2} h f^2 \nabla b \\
 & \quad + \beta (\mathbf{u} \cdot (\nabla g - \frac{1}{2} h \nabla f)) \nabla b - \frac{1}{2} h f \nabla g + \nabla (ghf) \\
 & = \frac{1}{2h} \nabla (h^2 \mathbf{u} \cdot \nabla g) - \frac{\beta}{2} h (\mathbf{u} \cdot \nabla f) \nabla b + \frac{\beta}{2} h f^2 \nabla b + \beta (\mathbf{u} \cdot \nabla g) \nabla b \\
 & = \mathcal{Q}_b[h, \beta b, \mathbf{u}].
 \end{aligned}$$

Thus the identity (6.1) holds, and the Proposition is proved. \square

Proof of Theorem 1.1. Let $N \geq 4$, $b \in \dot{H}^{N+2}$ and $(\zeta_0, \mathbf{u}_0) \in H^N \times X^N$ satisfying (1.7) with $h_*, h^* > 0$. Denote $h_0 = 1 + \epsilon \zeta_0 - \beta b$ and

$$\mathbf{v}_0 = h_0^{-1} \mathfrak{T}[h_0, \beta b] \mathbf{u}_0 = \mathbf{u}_0 + \mu \mathcal{T}[h_0, \beta b] \mathbf{u}_0,$$

where we recall that the operator \mathcal{T} is defined in (1.2). Using Lemma 2.1, 2.2 and 2.5, one easily checks that $\mathbf{v}_0 \in Y^N$, $\text{curl } \mathbf{v}_0 \in H^{N-1}$ and

$$|\mathbf{v}_0|_{Y^N} + |\text{curl } \mathbf{v}_0|_{H^{N-1}} \leq \mathbf{C}_0 (|\zeta|_{H^N} + |\mathbf{u}_0|_{X^N})$$

with $\mathbf{C}_0 = C(\mu, h_*^{-1}, h^*, \beta |\nabla b|_{H^{N+1}}, \epsilon |\zeta_0|_{H^N}, \epsilon |\mathbf{u}_0|_{X^N})$. One may thus apply Propositions 5.1 and 5.3: there exists $(\zeta, \mathbf{v}) \in C([0, T]; H^N \times Y^N)$ strong solution to (1.3); and one may restrict

$$T^{-1} = C(\mu, h_*^{-1}, h^*) F(\beta |\nabla b|_{H^{N+1}}, \epsilon |\zeta_0|_{H^N}, \epsilon |\mathbf{u}_0|_{X^N})$$

such that, for any $t \in [0, T]$, (1.7) holds with $\tilde{h}_* = h_*/2$, $\tilde{h}^* = 2h^*$, and

$$\mathcal{E}^N(\zeta, \mathbf{v}) + |\text{curl } \mathbf{v}|_{H^{N-1}}^2 \leq \mathbf{C}_0 (\mathcal{E}^N(\zeta_0, \mathbf{v}_0) + |\text{curl } \mathbf{v}_0|_{H^{N-1}}^2).$$

By Lemmata 2.1, 2.4 and 2.11 as well as Proposition 6.1, setting $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v})$ defines $(\zeta, \mathbf{u}) \in C([0, T]; H^N \times X^N)$ strong solution to (1.1),

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

and one has

$$\sup_{t \in [0, T]} (|\zeta|_{H^N}^2 + |\mathbf{u}|_{X^N}^2) \leq \mathbf{C}_0 (|\zeta_0|_{H^N}^2 + |\mathbf{u}_0|_{X^N}^2).$$

We thus constructed a strong solution to the Cauchy problem for (1.1) with initial data (ζ_0, \mathbf{u}_0) . The uniqueness of the solution follows from the uniqueness in Proposition 5.1 and Proposition 6.1. The continuity of the flow map follows from Proposition 5.3 and Lemmata 2.1, 2.4 and 2.11. This concludes the proof of Theorem 1.1.

Proof of Theorem 1.2. By the assumptions of Theorem 1.2, one has $\zeta_0 \in H^N$, $\nabla \psi_0 \in H^N$ and therefore, by Lemmata 2.1, 2.2 and 2.4, $\mathbf{u}_0 \in X^N$ and

$$|\mathbf{u}_0|_{X^N} \leq C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^{N-1}}, \epsilon |\zeta_0|_{H^N}) |\nabla \psi_0|_{H^N}.$$

Thus Theorem 1.1 applies, $(\zeta_{\text{GN}}, \mathbf{u}_{\text{GN}})$ is well-defined, and one can restrict T as in the Proposition to ensure that

$$\sup_{t \in [0, T]} (|\zeta_{\text{GN}}|_{H^N} + |\mathbf{u}_{\text{GN}}|_{X^N}) \leq \mathbf{C}_0,$$

with $\mathbf{C}_0 = C(\mu, h_\star^{-1}, h^\star, \beta |\nabla b|_{H^{N+1}}, \epsilon |\zeta_0|_{H^N}, \epsilon |\nabla \psi_0|_{H^N})$.

Now, slightly adapting the proof of [32, Prop. 5.8] and denoting

$$\mathbf{v}_{\text{ww}} \stackrel{\text{def}}{=} \nabla \psi_{\text{ww}}, \quad \mathbf{u}_{\text{ww}} \stackrel{\text{def}}{=} \mathfrak{I}[h_{\text{ww}}, \beta b]^{-1}(h_{\text{ww}} \mathbf{v}_{\text{ww}})$$

with $h_{\text{ww}} \stackrel{\text{def}}{=} 1 + \epsilon \zeta_{\text{ww}} - \beta b$, one finds that $(\zeta_{\text{ww}}, \mathbf{u}_{\text{ww}}) \in C(0, T; H^N \times X^N)$ satisfies (1.1) up to remainder terms $r_{\text{ww}}, \mathbf{r}_{\text{ww}}$, with

$$\sup_{t \in [0, T]} (|r_{\text{ww}}|_{H^{N-6}} + |\mathbf{r}_{\text{ww}}|_{H^{N-6}}) \leq \mu^2 \mathbf{C}_{\text{ww}},$$

with $\mathbf{C}_{\text{ww}} = C(\mu, h_\star^{-1}, \beta |b|_{H^N}, \epsilon |\zeta_{\text{ww}}|_{H^N}, \epsilon |\nabla \psi_{\text{ww}}|_{H^N})$. Proposition 6.1 immediately extends to non-trivial remainder terms, and it follows that $(\zeta_{\text{ww}}, \mathbf{v}_{\text{ww}}) \in C([0, T]; H^N \times Y^N)$ satisfies (1.3) up to the small remainder terms $r_{\text{ww}}, \mathbf{r}_{\text{ww}}$. We apply Proposition 5.2 and deduce that

$$(|\zeta_{\text{ww}} - \zeta_{\text{GN}}|_{H^{N-6}} + |\mathbf{v}_{\text{ww}} - \mathbf{v}_{\text{GN}}|_{Y^{N-6}})(t) \leq \mu^2 \mathbf{C} t,$$

with \mathbf{C} as in the statement. This concludes the proof of Theorem 1.2.

Appendix A. Energy estimates from the original formulation

As mentioned in the introduction, one can obtain energy estimates directly from system (1.1), rather than from system (1.3), as carried out in this work. We roughly sketch the different steps below.

Quasilinearization of the system. Let $(\zeta, \mathbf{u}) \in C([0, T]; H^{|\alpha|} \times X^{|\alpha|})$ satisfies (1.1), with α a non-zero multi-index and $T > 0$. Assume that b is sufficiently smooth, $|\alpha|$ is sufficiently large, and (1.7) holds. Then one can check that $\zeta_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \zeta$ and $\mathbf{u}_{(\alpha)} \stackrel{\text{def}}{=} \partial^\alpha \mathbf{u}$ satisfy

$$\begin{cases} \partial_t \zeta_{(\alpha)} + \epsilon \nabla \cdot (\mathbf{u} \zeta_{(\alpha)}) + \nabla \cdot (h \mathbf{u}_{(\alpha)}) = r_{(\alpha)}, \\ (\text{Id} + \mu \mathcal{T}[h, \beta b]) \partial_t \mathbf{u}_{(\alpha)} + \nabla \zeta_{(\alpha)} + \epsilon (\mathbf{u} \cdot \nabla) \mathbf{u}_{(\alpha)} + \mu \epsilon \mathcal{Q}_{(\alpha)}[h, \beta b, \mathbf{u}] \mathbf{u}_{(\alpha)} \\ \hspace{20em} = \mathbf{r}_{(\alpha)}, \end{cases} \quad (\text{A.1})$$

with $h = 1 + \epsilon \zeta - \beta b$ and (abusing notations)

$$\begin{aligned} \mathcal{Q}_{(\alpha)}[h, \beta b, \mathbf{u}] \mathbf{u}_{(\alpha)} &\stackrel{\text{def}}{=} \frac{-1}{3h} \nabla \left(h^3 ((\mathbf{u} \cdot \nabla)(\nabla \cdot \mathbf{u}_{(\alpha)})) \right) \\ &+ \frac{\beta}{2h} \left(\nabla (h^2 (\mathbf{u} \cdot \nabla)(\mathbf{u}_{(\alpha)} \cdot \nabla b)) - h^2 ((\mathbf{u} \cdot \nabla)(\nabla \cdot \mathbf{u}_{(\alpha)})) \nabla b \right) \\ &\hspace{10em} + \beta^2 ((\mathbf{u} \cdot \nabla)(\mathbf{u}_{(\alpha)} \cdot \nabla b)) \nabla b \end{aligned}$$

and where $(r_{(\alpha)}, \mathbf{r}_{(\alpha)}) \in C([0, T]; L^2 \times Y^0)$ satisfies

$$|r_{(\alpha)}|_{L^2} + |\mathbf{r}_{(\alpha)}|_{Y^0} \lesssim |\zeta|_{H^{|\alpha|}} + |\mathbf{u}|_{X^{|\alpha|}}. \quad (\text{A.2})$$

The system (A.1) satisfied by $(\zeta_{(\alpha)}, \mathbf{u}_{(\alpha)})$ is nothing but the linearized system (1.1) around (ζ, \mathbf{u}) , from which order-zero operators have been discarded. The estimate (A.2) would follow as in the proof of Proposition 3.1, and in particular using quasilinearization formulas derived in Section 2.

A priori energy estimates. For sufficiently smooth and finite-energy solutions of (A.1), we add the L^2 -inner product of the first equation with $\zeta_{(\alpha)}$ and the one of the second equation with $h \mathbf{u}_{(\alpha)}$. After some cancellations, integrations by parts and rearrangements, we find

$$\frac{d}{dt} \mathcal{F}_{(\alpha)} + \epsilon \mathcal{G}_{(\alpha)} = \int_{\mathbb{R}^d} r_{(\alpha)} \zeta_{(\alpha)} + h \mathbf{r}_{(\alpha)} \cdot \mathbf{u}_{(\alpha)} \, dx, \quad (\text{A.3})$$

where

$$\mathcal{F}_{(\alpha)} \stackrel{\text{def}}{=} \frac{1}{2} \int_{\mathbb{R}^d} \zeta_{(\alpha)}^2 + h|\mathbf{u}_{(\alpha)}|^2 + \mu h\mathcal{T}[h, \beta b]\mathbf{u}_{(\alpha)} \cdot \mathbf{u}_{(\alpha)} \, dx$$

and

$$\begin{aligned} \mathcal{G}_{(\alpha)} \stackrel{\text{def}}{=} & \frac{1}{2} \int_{\mathbb{R}^d} (\nabla \cdot \mathbf{u}) \zeta_{(\alpha)}^2 - (\partial_t \zeta + \nabla \cdot (h\mathbf{u})) |\mathbf{u}_{(\alpha)}|^2 \\ & - \frac{\mu}{3} (3h^2 \partial_t \zeta + \nabla \cdot (h^3 \mathbf{u})) (\nabla \cdot \mathbf{u}_{(\alpha)})^2 \\ & + \mu (2h \partial_t \zeta + \nabla \cdot (h^2 \mathbf{u})) (\beta \nabla b \cdot \mathbf{u}_{(\alpha)}) \nabla \cdot \mathbf{u}_{(\alpha)} \\ & - \mu (\partial_t \zeta + \nabla \cdot (h\mathbf{u})) (\beta \nabla b \cdot \mathbf{u}_{(\alpha)})^2 \, dx. \end{aligned}$$

By Lemma 2.3 and Cauchy-Schwarz inequality, we find that

$$|\zeta_{(\alpha)}|_{L^2}^2 + |\mathbf{u}_{(\alpha)}|_{X^0}^2 \lesssim \mathcal{F}_{(\alpha)} \quad \text{and} \quad \mathcal{G}_{(\alpha)} \lesssim |\zeta_{(\alpha)}|_{L^2}^2 + |\mathbf{u}_{(\alpha)}|_{X^0}^2.$$

Using (A.2) and again Cauchy-Schwarz inequality, Gronwall's Lemma to the differential equation (A.3) yields (locally in time) the control of the energy $\mathcal{F}_{(\alpha)}$. Proceeding as in Sections 4 and 5, one may then set up a Picard iteration scheme which yields the strong local well-posedness of the Cauchy problem for system (1.1).

Appendix B. Derivation of the Green-Naghdi system

Our work is based on a non-standard formulation of the Green-Naghdi system. We would like to motivate the relevance of this formulation (the verification of the equivalence between the different formulations is provided in Proposition 6.1). Below, we formally derive the non-standard formulation of the Green-Naghdi system from the Hamiltonian formulation of the water waves system, by approximating the associated Hamiltonian functional. This study, which was essentially provided in [12], has the advantage of revealing in a very straightforward way the Hamiltonian structure of the non-standard formulation of the Green-Naghdi system (and therefore the associated conservation laws) and giving a natural physical interpretation of the variables at stake.

Let us first recall the canonical Hamiltonian structure of the water waves system as brought to light by [54] and Craig-Sulem [19, 18]. Define the

following Hamiltonian functional

$$\mathcal{H}(\zeta, \psi) \stackrel{\text{def}}{=} \frac{1}{2} \int_{\mathbb{R}^d} \zeta^2 + \frac{1}{\mu} \psi G^\mu[\epsilon\zeta, \beta b] \psi \quad (\text{B.1})$$

where $\psi(t, X) = \phi(t, X, \epsilon\zeta(t, X))$ is the trace of the velocity potential at the surface, and G^μ is the Dirichlet-to-Neumann operator, defined by

$$G^\mu : \varphi \mapsto \sqrt{1 + \mu|\epsilon\nabla\zeta|^2}(\partial_n\phi) \Big|_{z=\epsilon\zeta} = (\partial_z\phi) \Big|_{z=\epsilon\zeta} - \mu(\epsilon\nabla\zeta) \cdot (\nabla_X\phi) \Big|_{z=\epsilon\zeta},$$

where ϕ is the unique solution (see *e.g.* [32] for a detailed and rigorous analysis) to

$$\begin{cases} \mu\Delta_X\phi + \partial_z^2\phi = 0 & \text{in } \{(X, z) \in \mathbb{R}^{d+1}, -1 + \beta b(X) \leq z \leq \epsilon\zeta(X)\}, \\ \phi(X, \epsilon\zeta(X)) = \varphi \\ (\partial_z\phi - \mu(\beta\nabla b) \cdot (\nabla_X\phi))(X, -1 + \beta b(X)) = 0. \end{cases} \quad (\text{B.2})$$

The operator G^μ is well-defined provided $h \stackrel{\text{def}}{=} 1 + \epsilon\zeta - \beta b \geq h_\star > 0$, and one can then show that the Zakharov/Craig-Sulem formulation of the water waves system simply reads

$$\partial_t \begin{pmatrix} \zeta \\ \psi \end{pmatrix} = \begin{pmatrix} 0 & \text{Id} \\ -\text{Id} & 0 \end{pmatrix} \begin{pmatrix} \delta_\zeta \mathcal{H} \\ \delta_\psi \mathcal{H} \end{pmatrix}. \quad (\text{B.3})$$

If one reformulates (in dimension $d = 2$) the above system using, instead of the canonical variables (ζ, ψ) , the variables ζ and $\mathbf{v} = (v_1, v_2)^\top \stackrel{\text{def}}{=} \nabla\psi$, then one obtains

$$\partial_t \begin{pmatrix} \zeta \\ v_1 \\ v_2 \end{pmatrix} = - \begin{pmatrix} 0 & \partial_1 & \partial_2 \\ \partial_1 & 0 & -q \\ \partial_2 & q & 0 \end{pmatrix} \begin{pmatrix} \delta_\zeta \mathcal{H} \\ \delta_{v_1} \mathcal{H} \\ \delta_{v_2} \mathcal{H} \end{pmatrix}. \quad (\text{B.4})$$

where $q = \frac{\text{curl}\mathbf{v}}{h}$. Of course, in our situation, $q \equiv 0$ since $\mathbf{v} = \nabla\psi$, but this contribution is kept for the analogy with the Euler or Saint-Venant Hamiltonian formalism ; see *e.g.* [49]. Keeping this contribution turns out to be necessary for comparing with the standard formulation of the Green-Naghdi system in the general setting; see Proposition 6.1.

Recall that one has the identity [32, Prop. 3.35]

$$\frac{1}{\mu} G^\mu[\epsilon\zeta, \beta b] \psi = -\nabla \cdot (h\mathbf{u}), \quad \mathbf{u} \stackrel{\text{def}}{=} \frac{1}{1 + \epsilon\zeta - \beta b} \int_{-1+\beta b}^{\epsilon\zeta} \nabla_X \phi(\cdot, z) \, dz. \quad (\text{B.5})$$

so that the first equation in (B.3) or (B.4) simply reads

$$\partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0,$$

which is the first equation of the Green-Naghdi system. The system is then completed by constructing an evolution equation for \mathbf{u} , containing only differential operators, which is approximately satisfied by exact solutions of the water waves system, through asymptotic expansions with respect to the parameter $\mu \rightarrow 0$. This equation has different equivalent formulations in the literature as the equations have been rediscovered several times; in this paper we use (1.1), originating from [33, (26)] and justified in the sense of consistency in [32, Prop. 5.8].

Our strategy here is different: we obtain equations written with the original variables ζ, ψ (or ζ, \mathbf{v}) by using an asymptotic expansion of the Hamiltonian functional \mathcal{H} , and plugging it in (B.3) or (B.4). The strategy of deriving the Green-Naghdi system using an approximate Hamiltonian functional or Lagrangian is not new: it was already used in particular in [53, 17] (leading however to an ill-posed system of Green-Naghdi type) and in [40] to derive the Green-Naghdi system; see also [12, 31, 15, 16].

Let us recall the Dirichlet-to-Neumann expansion [32, Remark 3.39]

$$\frac{1}{\mu} G^\mu[\epsilon\zeta, \beta b]\psi = -\nabla \cdot (h\nabla\psi) + \mu\nabla \cdot (h\mathcal{T}[h, \beta b]\nabla\psi) + \mathcal{O}(\mu^2) \quad (\text{B.6})$$

with the notation

$$\mathcal{T}[h, b]V \stackrel{\text{def}}{=} \frac{-1}{3h} \nabla(h^3\nabla \cdot V) + \frac{1}{2h} \left(\nabla(h^2\nabla b \cdot V) - h^2\nabla b\nabla \cdot V \right) + \nabla b(\nabla b \cdot V).$$

It would therefore be natural to consider the approximate Hamiltonian functional from (B.1)

$$\mathcal{H}(\zeta, \psi) \approx \frac{1}{2} \int_{\mathbb{R}^d} \zeta^2 + \psi \left(-\nabla \cdot (h\nabla\psi) + \mu\nabla \cdot (h\mathcal{T}[h, \beta b]\nabla\psi) \right).$$

However, plugging this approximation into (B.3) or (B.4) yields an ill-posed system (in the sense that the linearized system around the trivial solution $\zeta = 0, \psi = 0$, in the flat-bottom case, exhibits unstable modes whose amplitude grows exponentially and arbitrarily rapidly for large frequencies). It is interesting to note that the obtained system corresponds to the one exhibited in [53, (10)-(11)] and [17, (14)-(15)] (in the one-dimension and flat-bottom situation) and, as pointed out in [40, (1.8a),(1.8b)], it reduces to a standard (also ill-posed) Boussinesq (or Kaup) system when the amplitude is small, that is withdrawing $\mathcal{O}(\mu\epsilon)$ terms.

The ill-posedness of the aforementioned systems can be anticipated from the fact that the approximate Hamiltonian functional is no longer positive, whereas the original one is; see [32, Prop. 3.9 and 3.12]. This issue can be avoided as follows: by (B.5) and (B.6), one has

$$\mathbf{u} = \nabla\psi - \mu\mathcal{T}[h, \beta b]\nabla\psi + \mathcal{O}(\mu^2), \quad \text{thus} \quad \nabla\psi = \mathbf{u} + \mu\mathcal{T}[h, \beta b]\mathbf{u} + \mathcal{O}(\mu^2).$$

Now, we notice that the operator $\mathfrak{T}[h, \beta b]$ defined by

$$\mathfrak{T}[h, \beta b]\mathbf{u} \stackrel{\text{def}}{=} h\mathbf{u} + \mu h\mathcal{T}[h, \beta b]\mathbf{u}$$

is a topological isomorphism (see Lemma 2.3), and therefore

$$\mathbf{u} = \mathfrak{T}[h, \beta b]^{-1}(h\nabla\psi) + \mathcal{O}(\mu^2). \quad (\text{B.7})$$

It is now natural to use the following approximation:

$$\begin{aligned} \mathcal{H}(\zeta, \psi) &= \frac{1}{2} \int_{\mathbb{R}^d} \zeta^2 + (\nabla\psi) \cdot (h\mathbf{u}) \\ &\approx \frac{1}{2} \int_{\mathbb{R}^d} \zeta^2 + (h\nabla\psi) \cdot \mathfrak{T}[h, \beta b]^{-1}(h\nabla\psi) \stackrel{\text{def}}{=} \mathcal{H}_{\text{GN}}(\zeta, \psi). \end{aligned} \quad (\text{B.8})$$

Now, plugging the new approximate Hamiltonian in (B.3) and (B.4) yields, respectively,

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ \partial_t \psi + \zeta + \frac{\epsilon}{2} |\mathbf{u}|^2 = \mu\epsilon(\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]), \end{cases} \quad (\text{B.9})$$

and

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ (\partial_t + \epsilon \mathbf{u}^\perp \text{curl})\mathbf{v} + \nabla \zeta + \frac{\epsilon}{2} \nabla(|\mathbf{u}|^2) = \mu\epsilon \nabla(\mathcal{R}[h, \mathbf{u}] + \mathcal{R}_b[h, \beta b, \mathbf{u}]), \end{cases} \quad (\text{B.10})$$

where we denote $\mathbf{v} = \nabla\psi$, $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\nabla\psi)$, $(u_1, u_2)^\perp \stackrel{\text{def}}{=} (-u_2, u_1)$, $\text{curl}(v_1, v_2) \stackrel{\text{def}}{=} \partial_1 v_2 - \partial_2 v_1$, and

$$\begin{aligned} \mathcal{R}[h, \mathbf{u}] &\stackrel{\text{def}}{=} \frac{\mathbf{u}}{3h} \cdot \nabla(h^3 \nabla \cdot \mathbf{u}) + \frac{1}{2} h^2 (\nabla \cdot \mathbf{u})^2, \\ \mathcal{R}_b[h, \beta b, \mathbf{u}] &\stackrel{\text{def}}{=} -\frac{1}{2} \left(\frac{\mathbf{u}}{h} \cdot \nabla(h^2(\beta \nabla b \cdot \mathbf{u})) + h(\beta \nabla b \cdot \mathbf{u}) \nabla \cdot \mathbf{u} + (\beta \nabla b \cdot \mathbf{u})^2 \right). \end{aligned}$$

System (B.10) is the system we study, and we show in Proposition 6.1 that it is equivalent to the standard formulation of the Green-Naghdi

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

system, namely (1.1). System (B.9) is immediately deduced in the situation $\text{curl } \mathbf{v} = 0$, and inherit the canonical Hamiltonian structure of the Zakharov/Craig-Sulem formulation of the water waves system. Notice that one can rewrite system (B.10) as

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ (\partial_t + \epsilon \mathbf{u}^\perp \text{curl}) \mathbf{v} + \nabla (\zeta + \epsilon \mathbf{u} \cdot \mathbf{v} - \frac{\epsilon}{2} \mathbf{u} \cdot \mathbf{u} - \frac{\epsilon \mu}{2} w^2) = 0, \end{cases} \quad (\text{B.11})$$

with $w = (\beta \nabla b) \cdot \mathbf{u} - h \nabla \cdot \mathbf{u}$. To our knowledge, system (B.11), as a new formulation for the Green-Naghdi system, has been first brought to light in [12, (4.3)-(4.4)] (in the flat bottom case, the formulation (B.9) appears in [40, (6.5)] and [46, (9.12)] but is quickly disregarded in favor of the aforementioned ill-posed model). It appears also in [31, (5.14)-(5.15)] (in the irrotational setting), [23, (30)] (in the flat bottom situation) and [39, (2.9)-(2.34)-(2.35)]. As a matter of fact, the latter references point out that system (B.11) echoes a formulation of the water waves system. Indeed, system (B.4) may be equivalently written as

$$\begin{cases} \partial_t \zeta + \nabla \cdot (h\mathbf{u}) = 0, \\ (\partial_t + \epsilon U^\perp \text{curl}) \mathbf{v} + \nabla (\zeta + \epsilon U \cdot \mathbf{v} - \frac{\epsilon}{2} U \cdot U - \frac{\epsilon \mu}{2} w^2) = 0, \end{cases} \quad (\text{B.12})$$

where $\mathbf{v}, \mathbf{u}, U, w$ are determined from the velocity potential, ϕ , by

$$\mathbf{v} = \nabla(\phi \Big|_{z=1+\epsilon\zeta}), \quad \mathbf{u} = \frac{1}{h} \int_{\beta b}^{1+\epsilon\zeta} \nabla \phi \, dz, \quad (U, w) = (\nabla_X \phi, \mu^{-1} \partial_z \phi) \Big|_{z=1+\epsilon\zeta}.$$

System (B.12) is determined by the sole variables ζ and \mathbf{v} (and b), after solving the Laplace problem (B.2). Now, by the identity (B.5) and chain rule, one has

$$U = \mathbf{v} - \mu \epsilon w \nabla \zeta \quad \text{and} \quad w = \epsilon U \cdot \nabla \zeta - \nabla \cdot (h\mathbf{u}).$$

It follows in particular

$$U = \mathbf{v} + \mathcal{O}(\mu) = \mathbf{u} + \mathcal{O}(\mu) \quad \text{and} \quad w = (\beta \nabla b) \cdot \mathbf{u} - h \nabla \cdot \mathbf{u} + \mathcal{O}(\mu),$$

and therefore (B.11) is immediately seen as a $\mathcal{O}(\mu^2)$ approximation of (B.12), with the abuse of notation $\mathbf{u} \stackrel{\text{def}}{=} \mathfrak{T}[h, \beta b]^{-1}(h\mathbf{v})$ and $w \stackrel{\text{def}}{=} (\beta \nabla b) \cdot \mathbf{u} - h \nabla \cdot \mathbf{u}$ being justified by the above approximations.

Bibliography

- [1] T. Alazard. A minicourse on the low Mach number limit. *Discrete Contin. Dyn. Syst. Ser. S*, 1(3):365–404, 2008.
- [2] T. Alazard and G. Métivier. Paralinearization of the Dirichlet to Neumann operator, and regularity of three-dimensional water waves. *Comm. Partial Differential Equations*, 34(12):1632–1704, 2009.
- [3] S. Alinhac. Existence d’ondes de raréfaction pour des systèmes quasi-linéaires hyperboliques multidimensionnels. *Comm. Partial Differential Equations*, 14(2):173–230, 1989.
- [4] B. Alvarez-Samaniego and D. Lannes. Large time existence for 3D water waves and asymptotics. *Invent. Math.*, 171(3):485–541, 2008.
- [5] B. Alvarez-Samaniego and D. Lannes. A Nash-Moser theorem for singular evolution equations. Application to the Serre and Green-Naghdi equations. *Indiana Univ. Math. J.*, 57(1):97–131, 2008.
- [6] E. Barthélemy. Nonlinear shallow water theories for coastal waves. *Surveys in Geophysics*, 25(3-4):315–337, 2004.
- [7] S. Bellec, M. Colin, and M. Ricchiuto. Discrete asymptotic equations for long wave propagation. *SIAM J. Numer. Anal.*, 54:3280–3299, 2016.
- [8] S. Benzoni-Gavage and D. Serre. *Multidimensional hyperbolic partial differential equations. First-order systems and applications*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2007.
- [9] J. L. Bona and R. Smith. The initial-value problem for the Korteweg-de Vries equation. *Philos. Trans. Roy. Soc. London Ser. A*, 278(1287):555–601, 1975.
- [10] P. Bonneton, F. Chazel, D. Lannes, F. Marche, and M. Tissier. A splitting approach for the fully nonlinear and weakly dispersive Green-Naghdi model. *J. Comput. Phys.*, 230(4):1479–1498, 2011.

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

- [11] D. Bresch and G. Métivier. Anelastic limits for Euler-type systems. *Appl. Math. Res. Express. AMRX*, (2):119–141, 2010.
- [12] R. Camassa, D. D. Holm, and C. D. Levermore. Long-time effects of bottom topography in shallow water. *Phys. D*, 98(2-4):258–286, 1996. Nonlinear phenomena in ocean dynamics (Los Alamos, NM, 1995).
- [13] R. Camassa, D. D. Holm, and C. D. Levermore. Long-time shallow-water equations with a varying bottom. *J. Fluid Mech.*, 349:173–189, 1997.
- [14] A. Castro and D. Lannes. Fully nonlinear long-waves models in presence of vorticity. *J. Fluid Mech.*, 759:642–675, 2014.
- [15] M. Chhay, D. Dutykh, and D. Clamond. On the multi-symplectic structure of the Serre-Green-Naghdi equations. *J. Phys. A*, 49(3):03LT01, 7, 2016.
- [16] D. Clamond, D. Dutykh, and D. Mitsotakis. Conservative modified Serre-Green-Naghdi equations with improved dispersion characteristics. *Commun. Nonlinear Sci. Numer. Simul.*, 45:245–257, 2017.
- [17] W. Craig and M. D. Groves. Hamiltonian long-wave approximations to the water waves problem. *Wave Motion*, 19(4):367–389, 1994.
- [18] W. Craig and C. Sulem. Numerical simulation of gravity waves. *J. Comput. Phys.*, 108(1):73–83, 1993.
- [19] W. Craig, C. Sulem, and P.-L. Sulem. Nonlinear modulation of gravity waves: a rigorous approach. *Nonlinearity*, 5(2):497–522, 1992.
- [20] V. Duchêne, S. Israwi, and R. Talhouk. A new class of two-layer Green-Naghdi systems with improved frequency dispersion. *Stud. Appl. Math.*, 137(3):356–415, 2016.
- [21] H. Fujiwara and T. Iguchi. A shallow water approximation for water waves over a moving bottom. *Advanced Studies in Pure Mathematics*, 64:77–88, 2014.

- [22] I. Gallagher. Résultats récents sur la limite incompressible. *Astérisque*, (299):Exp. No. 926, vii, 29–57, 2005. Séminaire Bourbaki. Vol. 2003/2004.
- [23] S. Gavriluk, H. Kalisch, and Z. Khorsand. A kinematic conservation law in free surface flow. *Nonlinearity*, 28(6):1805–1821, 2015.
- [24] S. Gavriluk and V. Teshukov. Generalized vorticity for bubbly liquid and dispersive shallow water equations *Contin. Mech. Thermodyn.*, 13:365–382, 2001.
- [25] A. E. Green and P. M. Naghdi. A derivation of equations for wave propagation in water of variable depth. *J. Fluid Mech.*, 78(02):237–246, 1976.
- [26] D. D. Holm. Hamiltonian structure for two-dimensional hydrodynamics with nonlinear dispersion. *Phys. Fluids*, 31(8):2371–2373, 1988.
- [27] T. Iguchi. A shallow water approximation for water waves. *J. Math. Kyoto Univ.*, 49(1):13–55, 2009.
- [28] S. Israwi. Derivation and analysis of a new 2D Green-Naghdi system. *Nonlinearity*, 23(11):2889–2904, 2010.
- [29] S. Israwi. Large time existence for 1D Green-Naghdi equations. *Nonlinear Analysis: Theory, Methods & Applications*, 74(1):81–93, 2011.
- [30] T. Kato. The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Arch. Ration. Mech. Anal.*, 58(3):181–205, 1975.
- [31] J. W. Kim, K. J. Bai, R. C. Ertekin, and W. C. Webster. A derivation of the Green-Naghdi equations for irrotational flows. *J. Engrg. Math.*, 40(1):17–42, 2001.
- [32] D. Lannes. *The water waves problem*, volume 188 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2013. Mathematical analysis and asymptotics.
- [33] D. Lannes and P. Bonneton. Derivation of asymptotic two-dimensional time-dependent equations for surface water waves propagation. *Physics of Fluids*, 21(1):016601, 2009.

WP OF GREEN-NAGHDI AND BOUSSINESQ-PEREGRINE

- [34] D. Lannes and F. Marche. A new class of fully nonlinear and weakly dispersive Green-Naghdi models for efficient 2D simulations. *J. Comput. Phys.*, 282:238–268, 2015.
- [35] O. Le Métayer, S. Gavrilyuk, and S. Hank. A numerical scheme for the Green-Naghdi model. *J. Comput. Phys.*, 229:2034–2045, 2010.
- [36] C. D. Levermore, M. Oliver, and E. S. Titi. Global well-posedness for models of shallow water in a basin with a varying bottom. *Indiana Univ. Math. J.*, 45(2):479–510, 1996.
- [37] Y. A. Li. Hamiltonian structure and linear stability of solitary waves of the Green-Naghdi equations. *J. Nonlinear Math. Phys.*, 9(suppl. 1):99–105, 2002.
- [38] Y. A. Li. A shallow-water approximation to the full water waves problem. *Comm. Pure Appl. Math.*, 59(9):1225–1285, 2006.
- [39] Y. Matsuno. Hamiltonian structure for two-dimensional extended Green-Naghdi equations. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 472(2190):20160127–, 2016.
- [40] J. Miles and R. Salmon. Weakly dispersive nonlinear gravity waves. *J. Fluid Mech.*, 157:519–531, 1985.
- [41] B. Mésognon-Gireau. The Cauchy problem on large time for the Water Waves equations with large topography variations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 34(1):89–118, 2017.
- [42] B. Mésognon-Gireau. The Cauchy problem on large time for a Boussinesq-Peregrine equation with large topography variations. *Adv. Differential Equations*, 22(7/8):457–504, 2017.
- [43] B. Mésognon-Gireau. The singular limit of the Water-Waves equations in the rigid lid regime. ArXiv preprint:1512.02424.
- [44] M. Oliver. Classical solutions for a generalized Euler equation in two dimensions. *J. Math. Anal. Appl.*, 215(2):471–484, 1997.
- [45] D. H. Peregrine. Long waves on a beach. *J. Fluid Mech.*, 27:815–827, 3 1967.

- [46] R. Salmon. Hamiltonian fluid mechanics. *Annual Review of Fluid Mechanics*, 20(1):225–256, 1988.
- [47] F. J. Seabra-Santos, D. P. Renouard, and A. M. Temperville. Numerical and experimental study of the transformation of a solitary wave over a shelf or isolated obstacle. *J. Fluid Mech.*, 176:117–134, 3 1987.
- [48] F. Serre. Contribution à l'étude des écoulements permanents et variables dans les canaux. *La Houille Blanche*, 6:830–872, 1953.
- [49] T. G. Shepherd. Symmetries, conservation laws, and Hamiltonian structure in geophysical fluid dynamics. *Advances in Geophysics*, 32:287–338, 1990.
- [50] P. Siritwat, C. Kaewmanee, and S. V. Meleshko. Symmetries of the hyperbolic shallow water equations and the Green-Naghdi model in lagrangian coordinates. *International Journal of Non-Linear Mechanics*, 2016.
- [51] C. H. Su and C. S. Gardner. Korteweg-de Vries equation and generalizations. III. Derivation of the Korteweg-de Vries equation and Burgers equation. *J. Mathematical Phys.*, 10:536–539, 1969.
- [52] M. E. Taylor. *Partial differential equations. III Nonlinear equations*, volume 117 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1997.
- [53] G. B. Whitham. Variational methods and applications to water waves. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 299, 06 1967.
- [54] V. E. Zakharov. Stability of periodic waves of finite amplitude on the surface of a deep fluid. *J. Appl. Mech. Tech. Phys.*, 9:190–194, 1968.

VINCENT DUCHÊNE
Univ. Rennes 1, CNRS,
IRMAR - UMR 6625,
F-35000 Rennes, France.
vincent.duchene@univ-rennes1.fr

SAMER ISRAWI
Mathématiques, Faculté des sciences I et
Ecole doctorale des sciences et
technologie, Université Libanaise,
Beyrouth, Liban.